

Analytic theory of the eight-vertex model

Vladimir V. Bazhanov¹ and Vladimir V. Mangazeev²

*Department of Theoretical Physics,
Research School of Physical Sciences and Engineering,
Australian National University, Canberra, ACT 0200, Australia.*

Abstract

We observe that the exactly solved *eight-vertex solid-on-solid model* contains an hitherto unnoticed arbitrary field parameter, similar to the horizontal field in the six-vertex model. The parameter is required to describe a continuous spectrum of the *unrestricted* solid-on-solid model, which has an infinite-dimensional space of states even for a finite lattice. The introduction of the continuous field parameter allows us to completely review the theory of functional relations in the eight-vertex/SOS-model from a uniform analytic point of view. We also present a number of analytic and numerical techniques for the analysis of the Bethe Ansatz equations. It turns out that different solutions of these equations can be obtained from each other by analytic continuation. In particular, for small lattices we explicitly demonstrate that the largest and smallest eigenvalues of the transfer matrix of the eight-vertex model are just different branches of the same multivalued function of the field parameter.

Keywords: eight-vertex model, Bethe Ansatz, Q-operator, functional relations

PACS numbers: 02.30.Ik, 05.30.-d, 11.25.Hf

¹email: Vladimir.Bazhanov@anu.edu.au

²email: vladimir@maths.anu.edu.au

1 Introduction

The powerful analytic and algebraic techniques discovered by Baxter in his pioneering papers [1–4] on the exact solution of the eight-vertex lattice model laid the foundation for many important applications in the theory of integrable systems of statistical mechanics and quantum field theory.

This paper concerns one of these techniques — the method of functional relations. Over the last three decades, since Baxter’s original works [1–4], this method has been substantially developed and applied to a large number of various solvable models. However, the status of this method in the eight-vertex model itself with an account of all subsequent developments has not been recently reviewed. This paper is intended to (partially) fill this gap. Here we will adopt an analytic approach exploiting the existence of an (hitherto unnoticed) continuous field parameter in the solvable eight-vertex solid-on-solid model of ref. [3].

For the purpose of the following discussion it will be useful to first summarize the key results of [1–4]. Here we will use essentially the same notations as those in [1]. Consider the homogeneous eight-vertex (8V) model on a square lattice of N columns, with periodic boundary conditions. The model contains three arbitrary parameters u , η and $\mathbf{q} = e^{i\pi\tau}$, $\text{Im } \tau > 0$, which enter the parametrization of the Boltzmann weights (the parameter $\mathbf{q}$ enters as the nome for the elliptic theta-functions). The parameters η and $\mathbf{q}$ are considered as constants and the spectral parameter u as a complex variable. We assume that the parameter η is real and positive, $0 < \eta < \pi/2$, which corresponds to the disordered regime [5] of the model.

The row-to-row transfer matrix of the model, $\mathbf{T}(u)$, possesses remarkable analytic properties. Any of its eigenvalues, $\mathsf{T}(u)$, is both (i) an entire function of the variable u , and (ii) satisfies Baxter’s famous functional equation,

$$\mathsf{T}(u) \mathsf{Q}(u) = \mathsf{f}(u - \eta) \mathsf{Q}(u + 2\eta) + \mathsf{f}(u + \eta) \mathsf{Q}(u - 2\eta), \quad (1.1)$$

where ¹

$$\mathsf{f}(u) = (\vartheta_4(u | \mathbf{q}))^N, \quad (1.2)$$

and $\mathsf{Q}(u)$ is an entire quasi-periodic function of u , such that

$$\mathsf{Q}(u + \pi) = \pm(-1)^{N/2} \mathsf{Q}(u), \quad \mathsf{Q}(u + 2\pi\tau) = \mathbf{q}^{-2N} e^{-2iuN} \mathsf{Q}(u). \quad (1.3)$$

These analytic properties completely determine all eigenvalues of the transfer matrix $\mathbf{T}(u)$. Indeed, Eq.(1.1) implies that the zeroes $u_1, u_2, \dots, u_n$, of $\mathsf{Q}(u)$ satisfy the Bethe Ansatz equations,

$$\frac{\mathsf{f}(u_k + \eta)}{\mathsf{f}(u_k - \eta)} = -\frac{\mathsf{Q}(u_k + 2\eta)}{\mathsf{Q}(u_k - 2\eta)}, \quad \mathsf{Q}(u_k) = 0, \quad k = 1, \dots, n. \quad (1.4)$$

These equations, together with the periodicity relations (1.3), define the entire function $\mathsf{Q}(u)$ (there will be many solutions corresponding to different eigenvectors). Once $\mathsf{Q}(u)$ is known the eigenvalue $\mathsf{T}(u)$ is evaluated from (1.1).

The entire functions $\mathsf{Q}(u)$ appearing in (1.1) are, in fact, eigenvalues of another matrix, $\mathbf{Q}(u)$, called the **Q**-matrix. Originally it was constructed [1] in terms of some special transfer matrices. A different, but related, construction of the **Q**-matrix was given in [2] and later on used in the book [5]. An alternative approach to the 8V-model was developed in [3, 4] where Baxter invented the “*eight-vertex*” *solid-on-solid* (SOS) model and solved it exactly by means

¹Here we use the standard theta-functions [6], $\vartheta_i(u | \mathbf{q})$, $i = 1, \dots, 4$, $\mathbf{q} = e^{i\pi\tau}$, $\text{Im } \tau > 0$, with the periods π and $\pi\tau$. Our spectral parameter u is shifted with respect to that in [1] by a half of the imaginary period, see Sect 2.5.1 for further details.

of the co-ordinate Bethe Ansatz. This approach provided another derivation of the same result (1.1)-(1.4), since the 8V-model is embedded within the SOS-model.

Baxter's $\mathbf{Q}$ -matrix (or the $\mathbf{Q}$ -operator) possesses various exceptional properties and plays an important role in many aspects of the theory of integrable systems. A complete theory of the $\mathbf{Q}$ -operator in the 8V-model is not yet developed. However for the simpler models related with the quantum affine algebra $U_q(\widehat{sl_2})$ (where the fundamental $\mathbf{L}$ -operators [7] are intertwined by the $\mathbf{R}$ -matrix of the six-vertex model) the properties of the $\mathbf{Q}$ -operator are very well understood [8]. In this case the $\mathbf{Q}$ -operators (actually, there are two different $\mathbf{Q}$ -operators, $\mathbf{Q}_+$ and $\mathbf{Q}_-$) are defined as traces of certain monodromy matrices associated with infinite-dimensional representations of the so-called q -oscillator algebra. The main algebraic properties of the $\mathbf{Q}$ -operators can be concisely expressed by a single factorization relation

$$\mathbf{T}_j^+(u) = \mathbf{Q}_+(u + (2j + 1)\eta)\mathbf{Q}_-(u - (2j + 1)\eta) \quad (1.5)$$

where $\mathbf{T}_j^+(u)$ is the transfer matrix associated with the infinite-dimensional highest weight representation of $U_q(sl_2)$ with an arbitrary (complex) weight $2j$. Remarkably, this relation alone leads to a simple derivation of *all* functional relations involving various “fusion” transfer matrices and $\mathbf{Q}$ -operators [8, 9]. For this reason Eq.(1.5) can be regarded as a *fundamental fusion relation*: once it is derived, no further algebraic work is required.

An important part of the theory of the $\mathbf{Q}$ -operators belongs to their analytic properties with respect to a certain parameter, which we call here the “field parameter”. In the context of conformal field theory (considered in [8, 9]) this is the “vacuum parameter”, which determines the Virasoro highest weight Δ ; in the six-vertex model it corresponds to the horizontal field. In fact, the very existence of two different solutions [8, 10] of the TQ-equation (1.1) can be simply illustrated by the fact that the spectrum of the transfer matrix does not depend on the sign of the field, whereas the spectrum of the $\mathbf{Q}$ -operator does.

It is well known that it is impossible to introduce an arbitrary field parameter into the “zero-field” or “symmetric” eight vertex model of [1] without destroying its integrability. However, such parameter is intrinsically present in the solvable SOS-model. It does not enter the Boltzmann weights, but arises from a proper definition of the space of states of the model. To realize this recall that the SOS-model [4] is an interaction-round-a-face model where the face variables ℓ_i (called the heights) take arbitrary integer values $-\infty < \ell_i < +\infty$. Its transfer matrix acts in an infinite-dimensional space of states even for a finite lattice. It has a continuous spectrum, parameterized by the eigenvalue of the operator which simultaneously increments all height variables, $\ell_i \rightarrow \ell_i + 1$, on the lattice. Indeed, taking into account the results of [11, 12], it is not difficult to conclude that the calculations of [4] require only a very simple modification to deduce that the eigenvalues of the SOS transfer matrix enjoy the same TQ-equation (1.1), but require different periodicity properties

$$\mathbf{Q}_\pm(u + \pi) = e^{\pm i\varphi} \mathbf{Q}_\pm(u), \quad \mathbf{Q}_\pm(u + 2\pi\tau) = q^{-2N} e^{\pm\psi} e^{-2iuN} \mathbf{Q}_\pm(u), \quad (1.6)$$

where the exponent φ is arbitrary. It is determined by the eigenvalue $\omega = e^{2i\eta\varphi/\pi}$ of the height translation operator² (the second exponent ψ is dependent on φ). It is natural to assume, that

²In [4] Baxter restricted the parameter η to the “rational” values $L\eta = m_1\pi + m_2\pi\tau$, $L, m_1, m_2 \in \mathbb{Z}$ and considered a finite-dimensional subspace of the whole space of states, regarding the values of heights *modulo* L . In this case the phase factors $\omega = e^{2i\eta\varphi/\pi}$ take quantized values $\omega^L = 1$ (see [11, 13] for further discussion of this point). Apart from providing the conceptual advantage of a finite-dimensional space of states, the above restriction on η and ω was not used anywhere else in [4] and, therefore, can be removed. The transfer matrix of the 8V-vertex model (reformulated as the SOS-model) acts only in the finite-dimensional subspace of the SOS space of states, corresponding to a discrete set of exponents $\varphi = k\pi$ and $\psi = 0$ (the value of N is assumed to be even).

the functions $Q_{\pm}(u)$ solving these equations, are eigenvalues of the Q -operators for the SOS-model. Of course, it would be very desirable to obtain their explicit definition (and generalize the algebraic result (1.5) to the SOS-model), however, many properties of these operators can already be deduced from the information about their eigenvalues.

In this paper we will develop the analytic theory of the functional relations for the SOS-model starting from the eigenvalue equations (1.1) and (1.6). Bearing in mind that the TQ-equation (1.1) arises from very non-trivial algebraic “fusion” relations [1], it is not surprising that it implies all other functional relations. The required calculations are essentially the same as those in [8, 9], apart from trivial modifications arising in the context of lattice models.

The eigenvalues $Q_{\pm}(u)$ are two linear independent “Bloch wave” solutions [8, 10] of the finite difference equation (1.1) for the same $T(u)$. Their quantum Wronskian $W(\varphi)$, defined as,

$$2iW(\varphi)f(u) = Q_+(u+\eta)Q_-(u-\eta) - Q_+(u-\eta)Q_-(u+\eta), \quad (1.7)$$

is a complicated function of φ , η and q , depending on the eigenvalue $T(u)$. The Bloch solutions $Q_{\pm}(u)$ are well defined provided the exponent φ does not take some “singular values” (see Eq.(2.17) below), where $W(\varphi)$ vanishes. Otherwise Eq.(1.1) has only one quasi-periodic solution, while the second linear independent solution does not possess any simple periodicity properties.

All singular cases (in fact, they split into different classes) can be effectively studied with a limiting procedure starting from a non-singular value of φ . In the simplest case, when η is generic and φ approaches the points $\varphi = k\pi$, $k \in \mathbb{Z}$, the solutions $Q_+(u)$ and $Q_-(u)$ smoothly approach the same value (which for even N coincides with the eigenvalue $Q(u)$ of the 8V-model).

A more complicated situation occurs when the field φ tends to a singular value, say $\varphi = 0$, while the parameter η simultaneously approaches some rational fraction of π , where the transfer matrix of the 8V-model has degenerate eigenvalues. The limiting value of $T(u)$ is always uniquely defined. However, if $T(u)$ is a degenerate eigenvalue, the limiting values of $Q_{\pm}(u)$ are not uniquely defined. They have “complete exact strings” of zeroes whose position can be made arbitrary by changing the direction of the two-parameter (η, φ) -limit. Obviously, this reflects a non-uniqueness of eigenvectors for degenerate states [13]. An immediate consequence of this phenomenon is that, for rational η , there is no unique algebraic definition of the Q -operator in the symmetric 8V-model. This explains an important observation of [14], that Baxter’s two Q -operators, constructed in [1] and [2], are actually different operators, with different eigenvalues for degenerate eigenstates.

Further, the eigenvalues $Q_{\pm}(u)$, considered as functions of φ , have rather complicated analytic properties. Besides having the (relatively simple) singular points discussed above, they are multivalued functions with algebraic branching points in the complex φ -plane. The corresponding multi-sheeted Riemann surface appears to be extremely complicated; we were only able to numerically explore it for some particular eigenvalues.

It is easy to see that the roots of the Bethe Ansatz equations (1.4), considered as functions of φ satisfy a system of the first order ordinary differential equations, $du_k/d\varphi = U_k(u_1, u_2, \dots, u_n)$, where U_k are meromorphic functions of their arguments. Using these equations one can analytically continue any particular solution of (1.4) along a continuous path between two points, corresponding to the same value of φ on different sheets of the Riemann surface. In general, the resulting set of roots $u_1, u_2, \dots, u_n$ differs from the initial one, but, of course, satisfies exactly the same equation (1.4). In other words, different solutions of the Bethe Ansatz equations and, therefore, different eigenvalues of the transfer matrix can be obtained from each other by the analytic continuation in the parameter φ .

Guided by the above observation one might be tempted to suggest that the Bethe Ansatz equations have *only one solution, considered as a function of φ* . Undoubtedly, that could be an

elegant resolution to the problem of completeness, which traditionally attracts a lot of attention in the literature (see, e.g., the recent papers [13,15] and references therein). At the moment we cannot prove or disprove the above assertion. The analytic structure of the eigenvalues of the eight-vertex SOS model is quite complicated and certainly deserves further detailed studies.

This material was planned as an introductory part for an extended version of our previous work [16,17] devoted to the connection of the 8V-model with the Painlevé transcendents. However, in the course of writing, we realized that a review of the theory of the functional relations in the 8V-model could be of interest to a much wider audience than originally intended and deserves a separate publication. A detailed account of the results presented in [16,17] will be given in the second paper of this series [18], which is totally devoted to the special $\eta = \pi/3$ case of the eight vertex model with an odd number of sites. There we will consider remarkable connections of this special model with various differential equations, including the celebrated Lamé, Mathieu, Painlevé III and Painlevé VI equations.

The organization of the paper is as follows. In Sect. 2 we present the analytic theory of the functional relations in the 8V/SOS-model. In Sect. 3 we discuss some applications of the quantum Wronskian relation. In particular, we show how it can be used for the analysis of the degenerate states. In Sect. 4 we completely classify eigenvalues of the transfer matrix of the symmetric 8V-model for small lattices of the size $N \leq 4$. We then study the analytic properties of the eigenvalues with respect to the field variable φ with a combination of analytic and numerical techniques. In Conclusion we briefly summarize obtained results. Basic properties of the 8V-model are reviewed in the Appendix.

2 Functional relations in the eight-vertex SOS model.

2.1 Overview

In this section we will outline the analytic theory of the functional relations in the SOS-model (which also covers the symmetric 8V-model). Actually most of the functional relations discussed below are quite universal and apply to a wider class of related model. They include the six-vertex model in a field [19], the restricted solid-on-solid (RSOS) model [20] and some integrable models of quantum field theory: the $c < 1$ conformal field theory [8] and the massive sine-Gordon model in a finite volume [21].

Let $T(u)$ and $Q(u)$ denote the eigenvalues of the transfer matrix and the Q -operator respectively and η is an arbitrary real parameter in the range

$$0 < \eta < \pi/2 . \quad (2.1)$$

In all subsequent derivations we will use only one general assumption about the properties of the eigenvalues:

We assume that $T(u)$ and $f(u)$ are entire periodic function of the variable u ,

$$T(u + \pi) = T(u), \quad f(u + \pi) = f(u) , \quad (2.2)$$

and that the function $Q(u)$ solving the TQ -equation,

$$T(u) Q(u) = f(u - \eta) Q(u + 2\eta) + f(u + \eta) Q(u - 2\eta) , \quad (2.3)$$

is also an entire (but not necessarily periodic) function of u .

For every particular model the above requirements are supplemented by additional, model-specific analyticity properties of $Q(u)$ (such as, for example, the imaginary period relation (1.3) for the 8V-model). These properties will only be used in Sect. 3 and 4; they are discussed at the end of this section.

As explained in the Introduction, once the additional analyticity properties are fixed, the functional equation (2.3) completely determines all eigenvalues $T(u)$ and $Q(u)$. For certain applications, however, it is more convenient to use other functional equations in addition to (or instead of) (2.3). We will show that all such additional functional relations in the SOS-model (and in the related models mentioned above) follow elementary from two ingredients:

- (i) the TQ-equation itself (Eqs. (2.2) and (2.3) above), and
- (ii) the fact that for the same eigenvalue $T(u)$ this equation has two different $[8, 10]$ linearly independent solutions for $Q(u)$ which are entire functions of u .

The only property of the function $f(u)$ essentially used in this Section is its periodicity (2.2). For technical reasons we will also assume that $f(u - \eta)$ and $f(u + \eta)$ do not have common zeroes. This is a very mild assumption, excluding rather exotic row-inhomogeneous models, which are beyond the scope of this paper.

2.2 General functional relations.

Since $T(u)$ is an entire function, Eq.(2.3) implies that the zeroes $u_1, u_2, \dots, u_n$ of any eigenvalue $Q(u)$ satisfy the same set of the Bethe Ansatz equations

$$\frac{f(u_k + \eta)}{f(u_k - \eta)} = -\frac{Q(u_k + 2\eta)}{Q(u_k - 2\eta)}, \quad Q(u_k) = 0, \quad k = 1, \dots, n, \quad (2.4)$$

where the number of zeroes, n , is determined by the model-specific analyticity properties.

For any given eigenvalue $T(u)$ introduce an infinite set of functions $T_k(u)$, $k = 3, 4, \dots, \infty$, defined by the recurrence relation

$$T_k(u + \eta) T_k(u - \eta) = f(u + k\eta) f(u - k\eta) + T_{k-1}(u) T_{k+1}(u), \quad k \geq 2, \quad (2.5)$$

where

$$T_0(u) \equiv 0, \quad T_1(u) \equiv f(u), \quad T_2(u) \equiv T(u). \quad (2.6)$$

This relation can be equivalently rewritten as

$$T(u) T_k(u + k\eta) = f(u - \eta) T_{k-1}(u + (k+1)\eta) + f(u + \eta) T_{k+1}(u + (k-1)\eta), \quad (2.7a)$$

or as

$$T(u) T_k(u - k\eta) = f(u + \eta) T_{k-1}(u - (k+1)\eta) + f(u - \eta) T_{k+1}(u - (k-1)\eta). \quad (2.7b)$$

Using the definition (2.5) one can easily express $T_k(u)$ in terms of $T(u)$ as a determinant

$$T_k(u) = (f^{(k)}(u))^{-1} \det \left\| M_{ab}(u + k\eta) \right\|_{1 \leq a, b \leq k-1}, \quad k \geq 2, \quad (2.8)$$

where the $(k-1)$ by $(k-1)$ matrix $M(u)_{ab}$, $1 \leq a, b \leq k-1$, is given by

$$M_{ab}(u) = \delta_{a,b} T(u - 2a\eta) - \delta_{a,b+1} f(u - (2a+1)\eta) - \delta_{a+1,b} f(u - (2a-1)\eta), \quad (2.9)$$

while the normalization factor reads

$$f^{(k)}(u) = \prod_{\ell=0}^{k-3} f(u - (k - 3 - 2\ell)\eta) . \quad (2.10)$$

Finally, expressing $T(u)$ from (2.3) through the corresponding eigenvalue $Q(u)$ one arrives to the formula

$$T_k(u) = Q(u - k\eta) Q(u + k\eta) \sum_{\ell=0}^{k-1} \frac{f(u + (2\ell - k + 1)\eta)}{Q(u + (2\ell - k)\eta) Q(u + (2\ell - k + 2)\eta)} , \quad (2.11)$$

valid for $k = 1, 2, \dots, \infty$. Note, that the Bethe Ansatz equations (2.4) guarantee that all the higher $T_k(u)$ with $k \geq 3$ are entire functions of u as well as $T(u)$. It is worth noting that these functions are actually eigenvalues of the “higher” transfer matrices, obtained through the algebraic fusion procedure [22]. In our analytic approach this information is, of course, lost. Nevertheless it will be useful to have in mind that the index k in the notation $T_k(u)$ refers to the dimension of the “auxiliary” space in the definition of the corresponding transfer matrix. Another convenient scheme of notation for higher transfer matrices (used, e.g., in [9]) is based on (half-)integer spin labels j , such that $k = 2j + 1$.

In a generic case Eq.(2.3) has two linear independent “Bloch wave” solutions $Q_{\pm}(u)$, defined by their quasi-periodicity properties,

$$Q_{\pm}(u + \pi) = e^{\pm i\varphi} Q_{\pm}(u), \quad (2.12)$$

where the exponent φ depends on the eigenvalue $T(u)$. These solutions satisfy the quantum Wronskian relation

$$2iW(\varphi)f(u) = Q_+(u + \eta)Q_-(u - \eta) - Q_+(u - \eta)Q_-(u + \eta) , \quad (2.13)$$

where $W(\varphi)$ does not depend on u . Indeed, equating the two alternative expressions for $T(u)$,

$$T(u)Q_+(u) = f(u - \eta)Q_+(u + 2\eta) + f(u + \eta)Q_+(u - 2\eta) , \quad (2.14)$$

and

$$T(u)Q_-(u) = f(u - \eta)Q_-(u + 2\eta) + f(u + \eta)Q_-(u - 2\eta) , \quad (2.15)$$

and writing $W(\varphi)$ as $W(\varphi|u)$ (to assume its possible u -dependence, which cannot be ruled out just from the definition (2.13)) one gets $W(\varphi|u + \eta) = W(\varphi|u - \eta)$. On the other hand, Eqs.(2.13), (2.2) and (2.12) imply a different periodicity relation $W(\varphi|v + \pi) = W(\varphi|v)$. For generic real η , these two periodicity relations can only be compatible if $W(\varphi|u)$ is independent of u ,

$$W(\varphi|u) \equiv W(\varphi) . \quad (2.16)$$

When η and φ are in general position, the eigenvalues $Q_{\pm}(u)$ are locally analytic functions of u , therefore, by continuity, Eq.(2.16) at generic φ holds also when η/π is a rational number. However, when φ takes special values (for example, in the symmetric 8-vertex model) Eq.(2.16) for rational η/π cannot be established by the analytic arguments only.

Obviously, the condition (2.12) defines $Q_{\pm}(u)$ up to arbitrary u -independent normalization factors. Using this freedom, it is convenient to assume the normalization³ such that neither

³In the context of the 8V/SOS-model this is the most natural normalization. The eigenvalues $Q_{\pm}(u)$ are factorized in products of theta functions (see (3.15) below) and the variation of φ only affects positions of zeroes. Obviously, the transfer matrix eigenvalues, $T(u)$, do not have any singularities in φ .

of $Q_{\pm}(u)$ vanishes identically (as a function of u) or diverges at any value of φ . Then the quantum Wronskian $W(\varphi)$ will take finite values, but still can vanish at certain isolated values of the exponent φ . These values are called *singular* in the sense that there is only one quasi-periodic solution (2.12), while the second linear independent solution of (2.3) does not possess the simple periodicity properties (2.12). As argued in [8], the singular exponents take values in the “dangerous” set

$$\varphi_{dang} = k\pi + \frac{\pi^2}{2\eta} \ell, \quad k, \ell \in \mathbb{Z}. \quad (2.17)$$

However, each eigenvalue has its own set of singular exponents, being a subset of (2.17).

Evidently, $Q(u)$ in (2.11) can be substituted by any of the two Bloch solutions $Q_{\pm}(u)$, so there are two alternative expressions for each $T_k(u)$. Further, multiplying (2.14) and (2.15) by $Q_-(u - 2\eta)$ and $Q_+(u - 2\eta)$ respectively, subtracting resulting equations and using (2.13) one obtains

$$2iW(\varphi)T(u) = Q_+(u + 2\eta)Q_-(u - 2\eta) - Q_+(u - 2\eta)Q_-(u + 2\eta). \quad (2.18)$$

The last result, combined with the determinant formula (2.8), gives

$$2iW(\varphi)T_k(u) = Q_+(u + k\eta)Q_-(u - k\eta) - Q_+(u - k\eta)Q_-(u + k\eta), \quad (2.19)$$

where $k = 0, 1, 2, \dots, \infty$.

All the functional relations presented above are general corollaries of the TQ-equation (2.2), (2.3).

2.3 Rational values of η

Let us now assume that

$$2L\eta = m\pi, \quad 1 \leq m \leq L - 1, \quad L \geq 2, \quad (2.20)$$

where m and L are mutually prime integers. Evidently,

$$2k\eta \neq 0 \pmod{\pi}, \quad 1 \leq k \leq L - 1. \quad (2.21)$$

Combining the expression (2.11) with (2.2) and (2.12), one immediately obtains the following functional relation,

$$T_{L+k}(u) = 2\cos(m\varphi)T_k(u + \frac{1}{2}m\pi) + T_{L-k}(u), \quad k = 1, 2, \dots \quad (2.22)$$

which shows that, for the rational η of the form (2.20), all higher $T_k(u)$ with $k \geq L$ are the linear combinations of a finite number of the lower $T_k(u)$ with $k \leq L$. This relation is a simple corollary of the TQ-equation. It always holds for the rational values of η and does not require the existence of the second Bloch solution in (2.12) (indeed, Eq.(2.22) is independent of the sign of φ). Setting $k = 1$ in (2.22) one obtains

$$T_{L+1}(u) = 2\cos(m\varphi)f(u + \frac{1}{2}m\pi) + T_{L-1}(u). \quad (2.23)$$

This allows one to bring Eq.(2.5) with $k = L$ to the form

$$T_L(u + \eta)T_L(u - \eta) = \left(f(u + \frac{1}{2}m\pi) + e^{im\varphi}T_{L-1}(u)\right)\left(f(u + \frac{1}{2}m\pi) + e^{-im\varphi}T_{L-1}(u)\right) \quad (2.24)$$

where the periodicity (2.2) of the function $f(u)$ was taken into account. Thus, for the rational η , the equations (2.5) with $k = 2, 3, \dots, L - 1$ together with Eq.(2.24) form a closed system

of functional equations for a set of $L - 1$ eigenvalues $\{\mathbb{T}(u), \mathbb{T}_3(u), \dots, \mathbb{T}_L(u)\}$. Given that all $\mathbb{T}_k(u)$ with $k \geq 3$ are recursively defined through $\mathbb{T}(u)$, this system of equation leads to a single equation involving $\mathbb{T}(u)$ only. Indeed, substituting the determinant formulae (2.8) into (2.23) one obtains

$$\det \left\| \overline{\mathbb{M}}_{ab}(u) \right\|_{1 \leq a, b \leq L} = 0, \quad (2.25)$$

where the L by L matrix reads

$$\overline{\mathbb{M}}_{ab}(u) = \mathbb{M}_{ab}(u) - \omega \delta_{a,1} \delta_{b,L} \mathbf{f}(u - 3\eta) - \omega^{-1} \delta_{a,L} \delta_{b,1} \mathbf{f}(u + \eta) \quad (2.26)$$

with $\mathbb{M}_{ab}(u)$ given by (2.9) and $\omega = e^{\pm im\varphi}$.

2.3.1 Non-zero quantum Wronskian

Continuing the consideration of the rational case (2.20), let us additionally assume that both quasi-periodic solutions (2.12) exist and that their quantum Wronskian (2.13) is non-zero. It is worth noting that the functions $\mathbb{Q}_{\pm}(u)$ in this case cannot contain *complete exact strings*. A complete exact string (or, simply, a *complete string*) is a ring of L zeroes $u_1, \dots, u_L$, where each consecutive zero differs from the previous one by 2η , closing over the period π ,

$$u_{k+1} = u_k + 2\eta, \quad k = 1, \dots, L, \quad u_{L+1} = u_1 \pmod{\pi}. \quad (2.27)$$

It is easy to see that any such string manifests itself as a factor in the RHS of (2.13), but not in its LHS (unless, of course, $\mathbb{W}(\varphi) = 0$).

It follows from (2.12) that

$$\mathbb{Q}_{\pm}(u + m\pi) = e^{\pm im\varphi} \mathbb{Q}_{\pm}(u). \quad (2.28)$$

Using (2.19) and (2.20) one easily obtains the two equivalent relations,

$$e^{+im\varphi} \mathbb{T}_k(u) + \mathbb{T}_{L-k}(u + \tfrac{1}{2}m\pi) = \mathbb{C}(\varphi) \mathbb{Q}_+(u + k\eta) \mathbb{Q}_-(u - k\eta), \quad (2.29a)$$

and

$$e^{-im\varphi} \mathbb{T}_k(u) + \mathbb{T}_{L-k}(u + \tfrac{1}{2}m\pi) = \mathbb{C}(\varphi) \mathbb{Q}_+(u - k\eta) \mathbb{Q}_-(u + k\eta), \quad (2.29b)$$

where $k = 0, 1, \dots, L$ and

$$\mathbb{C}(\varphi) = \frac{\sin(m\varphi)}{\mathbb{W}(\varphi)}. \quad (2.30)$$

In particular, for $k = 0$ one gets,

$$\mathbb{T}_L(u + \tfrac{1}{2}m\pi) = \mathbb{C}(\varphi) \mathbb{Q}_+(u) \mathbb{Q}_-(u). \quad (2.31)$$

Quote also one simple but useful⁴ consequence of (2.29),

$$\log \frac{\mathbb{Q}_+(u + k\eta)}{\mathbb{Q}_-(u + k\eta)} - \log \frac{\mathbb{Q}_+(u - k\eta)}{\mathbb{Q}_-(u - k\eta)} = \log \left(\frac{e^{+im\varphi} \mathbb{T}_k(u) + \mathbb{T}_{L-k}(u + \tfrac{1}{2}m\pi)}{e^{-im\varphi} \mathbb{T}_k(u) + \mathbb{T}_{L-k}(u + \tfrac{1}{2}m\pi)} \right) \quad (2.32)$$

This first-order finite difference equation relates the ratio $\mathbb{Q}_+/\mathbb{Q}_-$ with the eigenvalues of the (higher) transfer matrices.

⁴Namely this relation with $k = 1$ was used in [23] to show that for rational values of η the expression for the non-linear mobility for the quantum Brownian particle in a periodic potential obtained in [8] exactly coincide with that of [24] found from the thermodynamic Bethe Ansatz.

Introduce the meromorphic functions

$$\Psi_{\pm}(u) = e^{\pm im\varphi} \sum_{\ell=0}^{L-1} \frac{f(u + (2\ell + 1)\eta)}{Q_{\pm}(u + 2\ell\eta) Q_{\pm}(u + (2\ell + 2)\eta)}, \quad (2.33)$$

such that

$$\mathsf{T}_L(u + \tfrac{1}{2}m\pi) = \left(Q_+(u)\right)^2 \Psi_+(u) = \left(Q_-(u)\right)^2 \Psi_-(u). \quad (2.34)$$

With this definition all the relations (2.29) reduce to a single relation which again can be written in two equivalent forms

$$\Psi_+(u) = C(\varphi) \frac{Q_-(u)}{Q_+(u)}, \quad \Psi_-(u) = C(\varphi) \frac{Q_+(u)}{Q_-(u)}. \quad (2.35)$$

Obviously,

$$\Psi_+(u)\Psi_-(u) = \left(C(\varphi)\right)^2, \quad \frac{\Psi_+(u)}{\Psi_-(u)} = \left(\frac{Q_-(u)}{Q_+(u)}\right)^2. \quad (2.36)$$

2.3.2 The RSOS regime and its vicinity

Further reduction of the functional relation in the rational case (2.20) occurs for certain special values of the field from the set

$$m\varphi = (r + 1)\pi, \quad r = 0, 1, 2, \dots. \quad (2.37)$$

Consider the effect of varying φ in the relation (2.31). The eigenvalue $\mathsf{T}_L(u)$ in the LHS will remain finite, so as the eigenvalues $Q_{\pm}(u)$ in the RHS. The latter also do not vanish identically (as functions of u) at any value of φ (see the discussion of our normalization assumptions before (2.17) above). Therefore the coefficient $C(\varphi)$, defined in (2.30), is always finite. This means that in the rational case (2.20), the quantum Wronskian, $W(\varphi)$, can only vanish at zeroes of the numerator in (2.30). However, the converse is not true: $W(\varphi)$ does not necessarily vanish when $C(\varphi) = 0$. Here we are interested in this latter case where

$$C(\varphi) = 0, \quad W(\varphi) \neq 0 \quad (2.38)$$

with φ from the set (2.37). By definition we call it the *RSOS regime*. The relations (2.29) and (2.31) reduce to

$$\mathsf{T}_k(u) = (-1)^r \mathsf{T}_{L-k}(u + \tfrac{1}{2}m\pi), \quad k = 1, \dots, L-1, \quad (2.39a)$$

and

$$\mathsf{T}_L(u) = 0. \quad (2.39b)$$

All these relations can be written as a single relation (in two equivalent forms involving only $Q_+(u)$ or $Q_-(u)$ respectively),

$$\Psi_+(u) = \Psi_-(u) = 0, \quad (2.40)$$

with $\Psi_{\pm}(u)$ defined by (2.33).

The special “truncation” relations (2.39), exactly coincide with those appearing in the RSOS-model [20]. These were obtained [25, 26] by the algebraic fusion procedure [27] and hold for all eigenvalues of the RSOS model. The above analysis shows that *all eigenvalues of the RSOS model are non-singular*. The quantum Wronskian of the Bloch solutions (2.12) is always non-zero (otherwise the coefficient $C(\varphi)$ in (2.29) would not have vanished). For this reason the

solutions of the Bethe Ansatz equations for the RSOS model cannot contain complete strings. Since, as argued in [13] the complete strings are necessary attributes of degenerate states, one arrives to a rather non-trivial statement: *the spectrum of the transfer matrix in the RSOS model is non-degenerate.*

Consider now the vicinity of the RSOS regime, when η and φ are approaching their limiting values given by (2.20) and (2.37) respectively. Interestingly, one can express some η - and φ -derivatives

$$\partial_\eta \mathsf{T}_k(u) = \frac{\partial}{\partial \eta} \mathsf{T}_k(u|\eta, \varphi), \quad \partial_\varphi \mathsf{T}_k(u) = \frac{\partial}{\partial \varphi} \mathsf{T}_k(u|\eta, \varphi), \quad (2.41)$$

calculated at the “RSOS point”,

$$(\eta, \varphi) = (m\pi/2L, \pi(r+1)/m), \quad (2.42)$$

in terms of the corresponding values of $\mathsf{Q}_\pm(u)$ and their first order u -derivatives

$$\mathsf{Q}'_\pm(u) = \frac{\partial}{\partial u} \mathsf{Q}_\pm(u|\eta, \varphi). \quad (2.43)$$

Using (2.19) one obtains,

$$\begin{aligned} \partial_\eta \left[\mathsf{T}_k(u) - (-1)^r \mathsf{T}_{L-k}(u + m\pi/2) \right] + L \partial_v \mathsf{T}_k(u) &= \\ &= \frac{L}{i\mathsf{W}(\varphi)} \left[\mathsf{Q}'_+(u + k\eta) \mathsf{Q}_-(u - k\eta) - \mathsf{Q}_+(u - k\eta) \mathsf{Q}'_-(u + k\eta) \right], \\ \partial_\varphi \left[\mathsf{T}_k(u) - (-1)^r \mathsf{T}_{L-k}(u + m\pi/2) \right] &= \\ &= \frac{m}{2\mathsf{W}(\varphi)} \left[\mathsf{Q}_+(u + k\eta) \mathsf{Q}_-(u - k\eta) + \mathsf{Q}_+(u - k\eta) \mathsf{Q}_-(u + k\eta) \right], \end{aligned} \quad (2.44)$$

where the expressions in the RHS are calculated directly at the point (2.42). According to the definitions (2.6), $\mathsf{T}_0(u)$ and $\mathsf{T}_1(u)$ do not depend on η and φ at all, therefore, one can express η - and φ -derivatives of $\mathsf{T}_{L-1}(u)$ and $\mathsf{T}_L(u)$ at the RSOS point (2.42) in terms of the values $\mathsf{Q}_\pm(u)$ and $\mathsf{Q}'_\pm(u)$.

2.4 Zero field case

Consider now the zero field limit $\varphi = 0$. Let us return to the case of an irrational η/π where the spectrum of the transfer matrix is non-degenerate. The eigenvalues $\mathsf{Q}_\pm(u)$, corresponding to the same eigenstate smoothly approach the same value at $\varphi = 0$. Moreover, adjusting a φ -dependent normalization of $\mathsf{Q}_\pm(u)$ one can bring their small φ expansion to the form

$$\mathsf{Q}_\pm(u) = \mathsf{Q}_0(u) \mp \varphi \overline{\mathsf{Q}}_0(u)/2 + O(\varphi^2), \quad \varphi \rightarrow 0, \quad (2.45)$$

where

$$\mathsf{Q}_0(u) = \mathsf{Q}_\pm(u)|_{\varphi=0}, \quad \overline{\mathsf{Q}}_0(u) = -2 \frac{d\mathsf{Q}_+(u)}{d\varphi} \Big|_{\varphi=0} = 2 \frac{d\mathsf{Q}_-(u)}{d\varphi} \Big|_{\varphi=0}. \quad (2.46)$$

From (2.12) it follows that

$$\mathsf{Q}_0(u + \pi) = \mathsf{Q}_0(u), \quad \overline{\mathsf{Q}}_0(u + \pi) = \overline{\mathsf{Q}}_0(u) + 2i\mathsf{Q}_0(u). \quad (2.47)$$

It is easy to see that the quasi-periodic part of $\overline{\mathsf{Q}}_0(u)$ is totally determined by $\mathsf{Q}_0(u)$,

$$\overline{\mathsf{Q}}_0(u) = \frac{2iu}{\pi} \mathsf{Q}_0(u) + \overline{\mathsf{Q}}_0^{(per)}(u). \quad (2.48)$$

However, the periodic part

$$\overline{Q}_0^{(per)}(u + \pi) = \overline{Q}_0^{(per)}(u), \quad (2.49)$$

can only be determined up to an additive term proportional to $Q_0(u)$. Indeed, consider the effect of an inessential normalization transformation

$$Q_{\pm}(u) \rightarrow e^{\pm\alpha\varphi} Q_{\pm}(u), \quad (2.50)$$

where α is a constant. The value of $Q_0(u)$ remains unchanged while the periodic part of $\overline{Q}_0(u)$ transforms as

$$\overline{Q}_0^{(per)}(u) \rightarrow \overline{Q}_0^{(per)}(u) - 2\alpha Q_0(u). \quad (2.51)$$

The quantum Wronskian relation (2.13) reduces to

$$Q_0(u + \eta) \overline{Q}_0(u - \eta) - Q_0(u - \eta) \overline{Q}_0(u + \eta) = 2i \dot{W}(0) f(u), \quad (2.52)$$

where

$$W'(0) = \left. \frac{dW(\varphi)}{d\varphi} \right|_{\varphi=0}. \quad (2.53)$$

The expression (2.19) now becomes

$$2iW'(0)T_k(u) = Q_0(u + k\eta) \overline{Q}_0(u - k\eta) - Q_0(u - k\eta) \overline{Q}_0(u + k\eta). \quad (2.54)$$

It is easy to see that at $\varphi = 0$ the TQ-equation (2.3) is satisfied if $Q(u)$ there is replaced by either of $Q_0(u)$ or $\overline{Q}_0(u)$. The same remark applies to the more general equation (2.11).

The Bethe Ansatz equations (2.4) for the zeroes of $Q_0(u)$ are the standard equations [1] arising in the analysis of the symmetric 8V-model. Exactly the same equations also hold for the zeroes of $\overline{Q}_0(u)$, but their usefulness is very limited. Even though $\overline{Q}_0(u)$ is an entire function of u , it lacks the simple periodicity (cf. (2.47)) and, therefore, does not possess any convenient product representation. Moreover, the transformation (2.51) affects the position of zeros of $\overline{Q}_0(u)$, making them ambiguous. All this renders the Bethe Ansatz equations for $\overline{Q}_0(u)$ useless. Fortunately, these equations are not really required for determination of $\overline{Q}_0(u)$. Once the zeros of $Q_0(u)$ are known the function $\overline{Q}_0(u)$ is explicitly calculated from (2.52) (see (3.29) below).

Additional functional relations arise in the rational case (2.20). These relations are straightforward corollaries of (2.29), (2.31) and (2.35). For instance, Eq.(2.29) gives

$$T_k(u) + T_{L-k}(u + m\pi/2) = C(0) Q_0(u + k\eta) Q_0(u - k\eta), \quad \varphi = 0, \quad (2.55)$$

where

$$C(0) = m/W'(0). \quad (2.56)$$

All these relations (with different k) can be equivalently re-written as a single relation

$$\sum_{\ell=0}^{L-1} \frac{f(u + (2\ell + 1)\eta)}{Q_0(u + 2\ell\eta) Q_0(u + (2\ell + 2)\eta)} = C(0), \quad \varphi = 0, \quad (2.57)$$

which is the $\varphi = 0$ version of (2.35). Setting $k = 0$ in (2.55) one gets

$$T_L(u + m\pi/2) = C(0) (Q_0(u))^2, \quad \varphi = 0. \quad (2.58)$$

Thus, at $\varphi = 0$ the eigenvalue $T_L(u + m\pi/2)$ becomes a perfect square. It only has *double* zeroes, whose positions coincide with the zeroes of $Q_0(u)$.

As is well known, in the rational case (2.20) the transfer matrix of the 8V-model has a degenerate spectrum (for sufficiently large values of $N \geq 2L$). We would like to stress here that the above relations (2.55)–(2.58) hold only for *non-degenerate* states. Actually, the assumption made in the beginning of this subsection, that $Q_+(u)$ coincides with $Q_-(u)$ when $\varphi = 0$, is true only for non-degenerate states. Removing this assumption and taking $\varphi \rightarrow 0$ limit in (2.31), while keeping η fixed by (2.20), one obtains

$$T_L(u + m\pi/2) = C(0) \bar{Q}_+(u) \bar{Q}_-(u), \quad \bar{Q}_\pm = \lim_{\varphi \rightarrow 0} Q_\pm(u). \quad (2.59)$$

For a degenerate state the eigenvalues $\bar{Q}_+(u)$ and $\bar{Q}_-(u)$ can only differ by positions of *complete exact strings*. This ambiguity does not affect any transfer matrix eigenvalues $T_k(u)$, since the complete strings trivially cancel out from (2.11). In principle, the complete strings can take arbitrary positions, however, for $\bar{Q}_\pm(u)$ they take rather distinguished positions. Indeed, due to (2.59), the zeroes of $\bar{Q}_\pm(u)$ manifest themselves as zeroes of $T_L(u + m\pi/2)$ which are uniquely defined even for the degenerate states. From the above discussion it is clear that $T_L(u)$ has either double zeroes or complete strings of zeroes. Further analysis of the degenerate case is contained in Sect.3.2.

2.5 Particular models

So far our considerations were rather general and covered several related models at the same time. For each particular model, one needs to specify additional properties, namely, (i) the explicit form of the function $f(u)$ and (ii) detailed analytic properties of the eigenvalues $Q_\pm(u)$. In this Section we will do this for three different models: the 8V/SOS-model, the 6V-model and the $c < 1$ conformal field theory.

2.5.1 The symmetric eight-vertex model

The basic properties of the 8V-model are briefly reviewed in the Appendix A. Readers who are not well familiar with the subject will benefit from reading this Appendix prior to the rest of the paper. Our notations are slightly different from those in Baxter's original papers [1–4]. The variables q , η , v and ρ used therein (hereafter denoted as q_B , η_B , v_B and ρ_B) are related to our variables q , η , v and ρ as

$$q^2 = q_B = e^{-\pi K'_B/K_B}, \quad \eta = \frac{\pi \eta_B}{2K_B}, \quad v = \frac{\pi v_B}{2K_B}, \quad \rho = \rho_B, \quad (2.60)$$

where K_B and K'_B are the complete elliptic integrals associated to the nome q_B . Here we fix the normalization of the Boltzmann weights as

$$\rho = 2 \vartheta_2(0|q)^{-1} \vartheta_4(0|q^2)^{-1}, \quad (2.61)$$

where

$$\vartheta_i(u|q), \quad i = 1, \dots, 4, \quad q = e^{i\pi\tau}, \quad \text{Im } \tau > 0, \quad (2.62)$$

are the standard theta functions [6] with the periods π and $\pi\tau$.

We denote the transfer matrix $\mathbf{T}$ and the $\mathbf{Q}$ -matrix from [1, 2] as $\mathbf{T}^B(v)$ and $\mathbf{Q}^B(v)$, remembering that our variable v is related to v_B by (2.60). Below we often use a shifted spectral parameter

$$u = v - \pi\tau/2, \quad (2.63)$$

simply connected to the variable v in (2.60). We also consider the re-defined matrices

$$\mathbf{T}(u) = (-i\mathbf{q}^{-1/4})^N e^{ivN} \mathbf{T}^B(v), \quad \mathbf{Q}(u) = e^{ivN/2} \mathbf{Q}^B(v) \quad (2.64)$$

where N is the number of columns of the lattice. The eigenvalues $\mathbf{T}(u)$ and $\mathbf{Q}(u)$ of these new matrices enjoy the following periodicity properties

$$\mathbf{T}(u + \pi) = \mathbf{T}(u), \quad \mathbf{T}(u + \pi\tau) = (-\mathbf{q})^{-N} e^{-2iuN} \mathbf{T}(u), \quad (2.65)$$

and

$$\text{8V-model:} \quad \mathbf{Q}(u + \pi) = s e^{i\pi N/2} \mathbf{Q}(u), \quad \mathbf{Q}(u + 2\pi\tau) = \mathbf{q}^{-2N} e^{-2iNu} \mathbf{Q}(u). \quad (2.66)$$

Here the “quantum number” $s = \pm 1$, is the eigenvalue of the operator $\mathcal{S}$, defined in (A.12). This operator always commutes with $\mathbf{T}(u)$ and $\mathbf{Q}(u)$.

Baxter’s TQ-equation (Eq.(4.2) of [1] and Eq.(87) of [2]) now takes the form (2.3) with

$$\mathbf{f}(u) = (\vartheta_4(u|\mathbf{q}))^N. \quad (2.67)$$

The main reason for the above redefinitions is to bring the TQ-equation to the universal form (2.3), where $\mathbf{T}(u)$ and $\mathbf{f}(u)$ are periodic functions of u (see Eq.(2.2)) for an arbitrary, odd or even, number of sites, N . This also helps to facilitate the considerations of the scaling limit in our next paper [18].

Comparing the first equation in (2.66) with the periodicity of the Bloch solutions (2.12) one concludes that the exponents φ read

$$\varphi^{(8V)} = \begin{cases} 0 \pmod{\pi}, & N = \text{even} \\ \frac{\pi}{2} \pmod{\pi}, & N = \text{odd} \end{cases} \quad (2.68)$$

Thus, for an even N the exponents of the symmetric 8V-model, with the cyclic boundary conditions, always belong to the “dangerous” set (2.17). For an odd N the exponents (2.68) fall into this set only for certain rational values of η/π . A notable example is the case $\eta = \pi/3$, considered in [16–18].

The imaginary period relations in (2.65) and (2.66) certainly deserve a detailed consideration. First, note that in (2.66) we only stated the periodicity with respect to the *double* imaginary period $2\pi\tau$, which always holds in all cases when the 8V-model has been exactly solved⁵. Actually, this is a rather overcautious statement which can be easily specialized further. For the following discussion assume a generic (i.e., irrational) value of η/π . Then for even N the Bloch solutions (2.12) always coincide (just as in the zero-field case of Sect.2.4). For odd N there are always two linearly independent Bloch solutions for each eigenvalue $\mathbf{T}(u)$, one with $s = +1$ and one with $s = -1$ (remind that in this case each eigenvalue of the transfer matrix is double-degenerate [28]). The existence of the “imaginary” period imposes rather non-trivial restrictions on the properties of the eigenvalues. Indeed, the second relation in (2.65) immediately implies that the function

$$\tilde{\mathbf{Q}}(u) = r \mathbf{q}^{N/2} e^{iuN} \mathbf{Q}(u + \pi\tau) \quad (2.69)$$

⁵Ref. [1] applies to rational η and arbitrary values of N , while ref. [2] applies to arbitrary η and even values N . It is reasonable to assume that (2.66) holds in general, however, the case of an arbitrary η and an odd N has never been considered.

where r is a constant, satisfies the TQ-equation (2.3) as well as $Q(u)$. Further, if $Q(u)$ is a Bloch solution

$$Q(u + \pi) = e^{i\varphi} Q(u) \quad (2.70)$$

with some φ then $\tilde{Q}(u)$ is also such a solution with the exponent

$$\tilde{\varphi} = \varphi + N\pi \pmod{2\pi}. \quad (2.71)$$

Obviously, there are two options, either $\tilde{Q}(u)$ is proportional to $Q(u)$ or it is proportional to the other linearly independent Bloch solution with the negated exponent “ $-\varphi$ ”. The first option is realized for even N ,

$$\text{8V-model, } N \text{ even:} \quad Q(u + \pi\tau) = r q^{-N/2} e^{-iuN} Q(u), \quad (2.72)$$

The constant $r = \pm 1$ is then the eigenvalue of the spin-reversal operator $\mathcal{R}$ (cf. (A.37)). The second option requires the exponent φ to be a half-an-odd integer fraction of π , it is realized for odd N ,

$$\text{8V-model, } N \text{ odd:} \quad Q_{\pm}(u + \pi\tau) = q^{-N/2} e^{-iuN} Q_{\mp}(u). \quad (2.73)$$

The above relations (2.72) and (2.73) were derived for irrational values of η/π , however they also hold in the rational case (2.20), if no additional degeneracy of the eigenvalues of the transfer matrix occurs (apart from the one related with the spin-reversal symmetry for odd N). The functional relation (2.31) can be then written in the form

$$T_L(u + \tfrac{1}{2}m\pi) = A e^{iNu} Q_+(u) Q_+(u + \pi\tau) \quad (2.74)$$

where A is a constant. This relation is identical to the one conjectured in [14]⁶.

2.5.2 The solid-on-solid model

The main idea of this paper is to study deformations of the eigenvalues $T(u)$ and $Q(u)$ under continuous variations of the exponents φ from their discrete values (2.68). As explained in the Introduction the resulting eigenvalues correspond to the unrestricted SOS-model. We will therefore assume the more general periodicity relations (1.6) for the Bloch wave solutions $Q_{\pm}(u)$, which hold for both odd and even N ,

$$\text{SOS-model:} \quad Q_{\pm}(u + \pi) = e^{\pm i\varphi} Q_{\pm}(u), \quad Q_{\pm}(u + 2\pi\tau) = q^{-2N} e^{\pm\psi} e^{-2iuN} Q_{\pm}(u), \quad (2.75)$$

where the exponent φ is arbitrary. The second exponent ψ is not an independent parameter, it is determined by φ (see the discussion in Section 3.1.2 below).

The second relation in (2.75) can be further refined for even N

$$\text{SOS-model, } N \text{ even:} \quad Q_{\pm}(u + \pi\tau) = q^{-N/2} e^{\pm\psi/2} e^{-iuN} Q_{\pm}(u), \quad (2.76)$$

whereas the periodicity of $T(u)$ remains the same (2.65) as in the 8V-model. However, there is no a general SOS-model analog of (2.73), as it is specific to half-odd exponents only. As a result Eq.(2.65) is replaced with

$$\text{SOS-model, } N \text{ odd:} \quad T(u + \pi) = T(u), \quad T(u + 2\pi\tau) = q^{-4N} e^{-4iuN} T(u). \quad (2.77)$$

Strictly speaking the use of the term “SOS-model” here is justified for even N only [3]. Nonetheless, we will use this term to indicate arbitrary values of the field parameter φ in general.

⁶The conjecture of [14] also covers a special case of degenerate states for rational values of η , where the relation (2.73) holds for the eigenvalues of the $\mathbf{Q}$ -matrix of [1] for even N .

2.5.3 Six-vertex model in a horizontal field

The allowed vertex configurations of the six-vertex model form a subset of those shown in Fig.12. Namely, the Boltzmann weights ϕ_7 and ϕ_8 are equal to zero. The remaining six weights will be parameterized as

$$\begin{aligned}\phi_1 &= e^{+H-i\eta} a, & \phi_2 &= e^{-H-i\eta} a, & \phi_3 &= e^{+H+i\eta} b, \\ \phi_4 &= e^{-H+i\eta} b, & \phi_5 &= e^{iu-2i\eta} c, & \phi_6 &= e^{iu-2i\eta} c,\end{aligned}\tag{2.78}$$

where H stands for the horizontal field

$$a = h(u + \eta), \quad b = h(u - \eta), \quad c = h(2\eta), \quad h(u) = 1 - e^{2iu} . \tag{2.79}$$

The above parametrization is simply related to that given in Eq.(12) of [13] (where the vertical field V is set to zero). The TQ-equation (eq.(11) of [13]) takes the form (2.2), (2.3), where

$$f(u) = (h(u))^N . \tag{2.80}$$

The Bloch solutions (2.12), corresponding to the eigenvectors of the transfer matrix with n “up-spins”, can be written as

$$Q_{\pm}(u) = e^{\pm i\varphi u/\pi} A_{\pm}(e^{2iu}) , \tag{2.81}$$

where $A_+(x)$ and $A_-(x)$ are polynomials in x of the degrees n and $(N - n)$, respectively, and

$$\varphi = \frac{i\pi H N}{2\eta} + \frac{\pi}{2} (N - 2n) . \tag{2.82}$$

Introduce new variables⁷

$$x = e^{2iu}, \quad q = e^{2i\eta}, \quad z = e^{2i\eta\varphi/\pi} . \tag{2.83}$$

Regarding x as a new spectral parameter instead of u and writing $T(u)$ and $f(u)$ as $T(x)$ and $f(x)$, respectively, one can rewrite (2.3) in the form

$$T(x) A_{\pm}(x) = z^{\pm 1} f(q^{-1} x) A_{\pm}(q^2 x) + z^{\mp 1} f(q x) A_{\pm}(q^{-2} x) , \tag{2.84}$$

where the polynomials $A_{\pm}(x)$ are defined in (2.81). This form is particularly convenient for the 6V-model.

2.5.4 Conformal field theory

The continuous quantum field theory version of Baxter’s commuting transfer matrices of the lattice theory was developed in [8,9,29]. These papers were devoted to the $c < 1$ conformal field theory (CFT). The parameters β and p used there define the central charge c and the Virasoro highest weight Δ ,

$$c = 1 - 6(\beta - \beta^{-1})^2, \quad \Delta = \left(\frac{p}{\beta}\right)^2 + \frac{c-1}{24} . \tag{2.85}$$

They are related to our η and φ as

$$2\eta = \beta^2 \pi, \quad \varphi = 2\pi p / \beta^2 . \tag{2.86}$$

⁷The parameter q should not be confused with the nome q in the 8V-model

The multiplicative spectral parameter λ used in those papers is related to our variable u as

$$\lambda^2 = -e^{-2iu} . \quad (2.87)$$

The eigenvalues of the CFT $\mathbf{Q}$ -operators $\mathbf{Q}_\pm(u)$ are entire functions of the variable u , satisfying the periodicity relation (2.12). Their leading asymptotics at large positive imaginary u read

$$\log \mathbf{Q}_\pm(u) = \frac{A}{\cos(\frac{\pi\eta}{\pi-2\eta})} e^{i\pi u/(\pi-2\eta)} + O(1), \quad u \rightarrow +i\infty, \quad |\operatorname{Re} u| < \pi/2 , \quad (2.88)$$

where A is a known constant [8]. Here we assumed that η does not belong to the set

$$\eta = \frac{\pi}{2} \left(1 - \frac{1}{2k}\right), \quad k = 1, 2, \dots, \infty . \quad (2.89)$$

At these special values of η the theory contains logarithmic divergences and the asymptotics (2.88) should be replaced with

$$\log \mathbf{Q}_\pm(u) = 2i(-1)^k A u e^{2iuk/\pi} + \mathbb{C} e^{2iuk} + O(1), \quad u \rightarrow +i\infty, \quad |\operatorname{Re} u| < \pi/2 , \quad (2.90)$$

where $\mathbb{C}$ is a regularization-dependent constant. The factorization formulae read⁸

$$\mathbf{Q}_\pm(u) = e^{\pm iu\varphi/\pi} \mathbf{A}_\pm(u), \quad \mathbf{A}_\pm(u) = \prod_{k=1}^{\infty} (1 - e^{-2i(u-u_k^\pm)}) , \quad (2.91)$$

where the zeroes $u_1^\pm, u_2^\pm, \dots$ accumulate at infinity along the straight line

$$u = \pi/2 + iy, \quad y \rightarrow +\infty . \quad (2.92)$$

Finally, the function $\mathbf{f}(u)$ in the case of CFT should be set to one⁹,

$$\mathbf{f}(u) \equiv 1 . \quad (2.93)$$

With these specializations the functional relations given above become identical to those previously obtained in [8, 9, 29].

2.6 Related developments and bibliography

The literature on the functional relation in solvable models is huge; therefore it would not be practical to mention all papers in the area. Our brief review is restricted only to a subset of publications directly related to the eight-vertex/six-vertex models and associated models of quantum field theory.

2.6.1 Transfer matrix relations

In the above presentation the entire functions $\mathbf{T}_k(u)$ with $k \geq 3$ were defined by the recurrence relation (2.5), which allows one to express them solely in terms of $\mathbf{T}(u)$, as in (2.8). No other additional properties of $\mathbf{T}_k(u)$ were used. However, as is well known, these functions are eigenvalues of the higher transfer matrices, usually associated with the so-called fusion procedure.

⁸Here we assumed that $0 < \eta < \pi/4$. When $\pi/4 < \eta < \pi/2$ the product in (2.91) should contain the standard Weierstrass regularization factors [8].

⁹Again, we have assumed that η does not fall into the set (2.89), otherwise $\mathbf{f}(u) = \exp(4A\eta e^{2iuk}/\pi)$.

This algebraic procedure provides a derivation of the functional relations for the higher transfer matrices based on decomposition properties of products of representations of the affine quantum groups. Originally, all these “transfer matrix relations” were obtained essentially in this way. We would like to stress that the logic of these developments was exactly opposite to that employed in our review. The goal was to find new techniques, independent of the TQ-relation, rather than to deduce everything from the latter. The first important contribution was made by Stroganov [30]. He gave an algebraic derivation of the first nontrivial relation in (2.5) (with $k = 2$),

$$T(u + \eta) T(u - \eta) - f(u + 2\eta) f(u - 2\eta) = O((u - u_0)^N) \quad (2.94)$$

in the vicinity of the point $u = u_0$ where the transfer matrices $T(u_0 + \eta)$ and $T(u_0 - \eta)$ become shift operators. Remarkably, this single relation alone contains almost all information about the eigenvalues $T(u)$. To illustrate this point consider, for instance, the 6V-model. For a chain of the length N each eigenvalue $T(u)$ is a trigonometric polynomial of the degree N , determined by $N + 1$ unknown coefficients. The mere fact that the LHS of (2.94) has an N -th order zero immediately gives N algebraic equations for these unknowns. Similar arguments, obviously, apply to the 8V-model. One additional equation is usually easy to find from some elementary considerations (e.g., from the large u asymptotics in the 6V-model). Further, in the thermodynamic limit, $N \rightarrow \infty$ with u kept fixed, Eq.(2.94) becomes a closed functional relation for the eigenvalues (its RHS vanishes). This is the famous “inversion relation” [30–32]. With additional analyticity assumptions it can be effectively used to calculate the eigenvalues of the transfer matrix at $N = \infty$. Recently, Eq.(2.94) was used to derive a new non-linear integral equation [33], especially suited for the analysis of high-temperature properties of lattice models.

Soon after [30] Stroganov derived [34] a particular case of (2.23) for the 6-vertex model with $\eta = \pi/6$ (i.e., for $L = 3$ and $m = 1$). He also found that for the case of an odd number of sites¹⁰ Eq.(2.25) takes the form

$$T(u - 2\eta) T(u) T(u + 2\eta) = f(u)f(u + 2\eta) T(u - 2\eta) + f(u - 2\eta)f(u + 2\eta) T(u) + f(u - 2\eta)f(u) T(u + 2\eta) . \quad (2.95)$$

He then used this equation to obtain Bethe Ansatz type equations for the zeroes of $T(u)$ and to reproduce Lieb’s celebrated result [35] for the residual entropy of the two-dimensional ice. Unfortunately, these results were left unpublished.

The ideas of [30, 34] were further developed in the *analytic Bethe Ansatz* [36] where the TQ-equation (or an analogous equation) is used essentially as a formal substitution to solve the transfer matrix functional equations. The notion of “higher” or “fused” R-matrices was developed in [22] from the point of view of representation theory. These R-matrices were calculated in [37] for the 6V-model, in [38–41] for the 8V-model and in [27] for the SOS-model. The functional relations (2.7) were given in [37] for the 6V-model and in [26] for 8V/SOS-model. The determinant identity (2.25) was discussed in [26, 42]. An algebraic derivation of the truncation relations (2.39) for the RSOS model [20] was given in [26]. A particular case of this truncation for the hard hexagon model [43] was previously discovered in [25]. An algebraic derivation of (2.23) in the zero-field six-vertex model is given in [44]. The idea of calculation of φ - and η -derivatives (2.44) at the RSOS point given in Sect.2.3.2 is borrowed from [45] and [46].

Remarkably, the same functional equations (2.5) (along with all their specializations in the rational case) arise in a related, but different context of the thermodynamic Bethe Ansatz [47]; see [48] for its application to the 8V-model. Usually this approach in lattice models is associated with non-linear integral equations. Here we refer to the functional form of these equations

¹⁰In our notations this corresponds to $\varphi = \pi/2 \pmod{\pi}$.

discovered in [49]. Further discussion of the correspondence of the functional relation method with the thermodynamic Bethe Ansatz and its generalizations for excited states can be found in [21, 29, 50–53].

2.6.2 Q-matrix and TQ-relations

As noted before, a full algebraic theory of the $\mathbf{Q}$ -matrix in the 8V-model is not yet developed. The idea of the construction of the $\mathbf{Q}$ -matrix in terms of some special transfer matrices belongs to Baxter. It is a key element of his original solution of the 8V-model. Readers interested in details should familiarize themselves with the Appendix C of [1] (along with other four appendices and, of course, the main text of that paper, which contain a wealth of important information on the subject). The results of [1] only apply for certain rational values of η . The construction of [1] and the set of allowed values of η were recently revised in [14]. A different construction for the $\mathbf{Q}$ -matrix, which works for an arbitrary η , was given [3]. Some remarks on a comparison between the two $\mathbf{Q}$ -matrices are given in Appendix A.

There are many related solvable models connected with the R-matrix of the 8V-model but having different $\mathbf{L}$ -operators and different *quantum spaces*. The general structure of the functional relations in all such models remains the same. In particular, they all possess a TQ-relation (though it may contain different scalar factors and require different analytic properties of the eigenvalues). In [3] Baxter also presented an extremely simple explicit formula for the matrix elements of the $\mathbf{Q}$ -matrix for the zero-field 6V-model in the sector with $N/2$ “up-spins” (the half filling). However, no such expression is known for the 8V-model, or the other sectors of the 6V-model. The quantum space of the 6V-model is build from the two-dimensional highest weight representation of $U_q(sl_2)$ at every site of the lattice. Curiously enough, if this representation is replaced with the general cyclic representation (arising at roots of unity, $q^L = 1$) then all matrix elements of the $\mathbf{Q}$ -matrix can be explicitly calculated [54] as a simple product involving only a two-spin interaction¹¹. Remarkably, the resulting $\mathbf{Q}$ -matrix exactly coincides with the transfer matrix of the chiral Potts model [55–57]; this allows one to view the latter as a “descendant of the six-vertex model” [54]. The generalization of this construction to the eight-vertex and the Kashiwara-Miwa model [58] is considered in [59]. Further developments of the theory of the $\mathbf{Q}$ -matrix and related topics (along with many important applications to various solvable models) can be found in [60–69].

Baxter’s original idea of the construction of $\mathbf{Q}$ -operators which utilizes traces of certain monodromy matrices was extended in [8, 9] for trigonometric models related with the quantum affine algebra $U_q(\widehat{sl}(2))$. It turned out that in the trigonometric case the situation is considerably simpler than for the 8-vertex model and the $\mathbf{Q}$ -operators coincide with some special transfer matrices. The corresponding $\mathbf{L}$ -operators are obtained as specializations of the universal $\mathbf{R}$ -matrix [70] to infinite-dimensional representations of the q -oscillator algebra in the “auxiliary space”. Although the calculations of [8, 9] were specific to the continuous quantum field theory, the same procedure can readily be applied to lattice models (see, e.g., [71–74] for the corresponding results for the 6V-model). In the case of the 6V-model with non-zero horizontal field this construction leads to two $\mathbf{Q}$ -matrices¹², whose eigenvalues $Q_{\pm}$ are precisely the “Bloch wave” solutions of the TQ-equation.

Note that functional relations which involves bi-linear combinations of $Q_{\pm}$, namely (2.19),

¹¹The factorization of the matrix elements of the $\mathbf{Q}$ -matrix is typical for quantum space representations without highest and lowest weights.

¹²As noted in [9], for the “half-filled” sector of the zero-field 6V-model these $\mathbf{Q}$ -matrices reduce to Baxter’s expression [3] mentioned above, as they, of course, should.

(2.29) and (2.31) are universal in the sense that they do not involve the model-specific function $f(u)$. These relations were derived in [8, 9] in the context of the conformal field theory. Similar relations previously appeared in the chiral Potts model [54, 75, 76], though the correspondence is not exact because there is no an additive spectral parameter in that model. Eq.(2.19) in the eight-vertex and the XXX-models was found in [10] and [77]. A special case (2.74) of the relation (2.31) involving Baxter's original $\mathbf{Q}$ -matrix [1] for the 8V-model was conjectured in [14]. Another special (zero-field) case (2.55) of the same relation (2.31) in conformal field theory was conjectured in [52].

3 Quantum Wronskian relation and singular eigenvalues

The quantum Wronskian relation (2.13), naturally arising in the above analysis of the TQ equation, is a very non-trivial functional relation. In this Section we show how this relation can be used for the analysis of the eigenvalues. In particular, we consider a practical question of the calculation of $Q_-(u)$ from a known $Q_+(u)$. Next, we study certain singular eigenvalues in the zero-field limit.

3.1 Solving the quantum Wronskian relation

3.1.1 Six-vertex model (multiplicative spectral parameter)

Let us first consider the example of the 6V-model. It is convenient to present the polynomials $A_{\pm}(x)$, defined in (2.81) in a factorized form,

$$A_+(x) = \rho_+ \prod_{k=1}^n (1 - x/x_k^+) , \quad A_-(x) = \rho_- \prod_{k=1}^{N-n} (1 - x/x_k^-) , \quad (3.1)$$

and rewrite the quantum Wronskian relation (2.13) as

$$z A_+(xq) A_-(x/q) - z^{-1} A_+(x/q) A_-(xq) = 2i W(z) f(x) . \quad (3.2)$$

This single functional equation determines both unknown polynomials $A_{\pm}(x)$, as well as the function $W(z)$, except for an obvious freedom to choose arbitrary x -independent factors $\rho_{\pm}$ without changing the normalization-independent combination

$$2i W(z)/(\rho_+ \rho_-) = z - z^{-1} , \quad (3.3)$$

fixed by Eq.(3.2) at $x = 0$. There will be many solutions to (3.2) corresponding to different eigenvectors of the transfer matrix. Indeed, setting there $x = q x_k^+$ and $x = q^{-1} x_k^+$, one obtains two relations

$$z A_-(x_k^+) A_+(x_k^+ q^2) = 2i W(z) f(x_k^+ q), \quad z^{-1} A_-(x_k^+) A_+(x_k^+ q^{-2}) = -2i W(z) f(x_k^+ q^{-1}) . \quad (3.4)$$

Taking their ratio, one comes back to the Bethe Ansatz equations for the zeroes $x_1^+, x_2^+, \dots, x_n^+$ of $A_+(x)$,

$$\frac{f(x_k^+ q)}{f(x_k^+ q^{-1})} = -z^2 \frac{A_+(x_k^+ q^2)}{A_+(x_k^+ q^{-2})}, \quad k = 1, 2, \dots, n \quad (3.5)$$

Similar equations (with z replaced by z^{-1}) hold for the zeroes $x_1^-, x_2^-, \dots, x_{N-n}^-$ of $A_-(x)$.

Given a solution of (3.5) (which defines $A_+(x)$) the corresponding polynomial $A_-(x)$ is uniquely determined by (3.2). For example, consider the case $N = 2n$. Comparing the leading power of x in (3.2) one gets

$$s \stackrel{\text{def}}{=} (x_1^+ x_2^+ \cdots x_n^+)^{-1} = x_1^- x_2^- \cdots x_n^- . \quad (3.6)$$

Let us rewrite (3.2) as a functional difference equation

$$z F(x/q) - z^{-1} F(xq) = \frac{f(x)}{A_+(xq)A_+(x/q)} \quad (3.7)$$

for the rational function

$$F(x) = \frac{1}{2iW(z)} \frac{A_-(x)}{A_+(x)} . \quad (3.8)$$

Performing the partial fraction decomposition of the RHS of (3.7) and sharing the poles between $F(xq)$ and $F(x/q)$ one obtains

$$\begin{aligned} F(x) &= \frac{1}{z(1-s^2)} \sum_{k=1}^n \frac{f(x_k^+ q)}{x_k^+ A_+(x_k^+ q^2) A'_+(x_k^+)} \left(\frac{x - s^2 x_k^+}{x - x_k^+} \right) \\ &= -\frac{z}{(1-s^2)} \sum_{k=1}^n \frac{f(x_k^+ q^{-1})}{x_k^+ A_+(x_k^+ q^{-2}) A'_+(x_k^+)} \left(\frac{x - s^2 x_k^+}{x - x_k^+} \right) \end{aligned} \quad (3.9)$$

where $A'_+(x)$ denotes the derivative of $A_+(x)$ with respect to x . The two alternative expressions given above are equivalent in virtue of the Bethe Ansatz equations (3.5). The formula (3.9) can be easily generalized to arbitrary values of n . We leave this as an exercise for the reader.

The above derivation has several limitations. Evidently the field parameter z should not take the values $z = \pm 1$, where the denominator in (3.8) vanishes. Further, the above sharing of the poles in (3.7) is only possible if all the roots x_k^+ are distinct, finite and non-zero. Moreover no pair of the roots could satisfy the condition $x_j^+/x_k^+ = q^{\pm 2}$. In all these cases the expression (3.9) becomes meaningless (typically the summand therein diverges).

The quantum Wronskian relation can also be solved directly for the coefficients of the polynomials (2.81)

$$A_+(x) = \sum_{k=0}^n a_k^+ x^k, \quad A_-(x) = \sum_{k=0}^{N-n} a_k^- x^k, \quad (3.10)$$

This approach lead to a system of algebraic equations which sometimes easier to analyze since it involves only symmetric functions of the Bethe roots, but not the roots themselves. As an example, consider the trivial eigenvalue

$$T(x) = z f(xq^{-1}) + z^{-1} f(xq) \quad (3.11)$$

corresponding to the ferromagnetic ground state (all spins down) of the six-vertex. One solution of (3.2) is obvious

$$A_+(x) = \rho_+ = 1 . \quad (3.12)$$

The finite difference equation (3.7) can be readily solved

$$\frac{A_-(x)}{2iW(z)} = -z \sum_{k=0}^N \binom{N}{k} \frac{(-xq)^k}{z^2 - q^{2k}} . \quad (3.13)$$

If z takes one the values $z = \pm q^k$, $k = 0, \dots, N$, the polynomial $A_-(x)$ reduces to a single power, x^k , and the corresponding Bloch solutions (2.81) become linearly dependent. These are the singular cases discussed in Sect.2.2. Indeed, choosing the normalization

$$\rho_- = \prod_{k=1}^N (z^2 - q^{2k}) \quad (3.14)$$

so that $A_-(x)$ remains finite and does not vanish identically for all z and taking into account the definition of z in (2.83), one can easily see from (3.3) and (3.12) that the singular exponents form a subset of (2.17), as expected.

3.1.2 Eight-vertex/SOS model

The formula similar to (3.9) holds for the 8-vertex/SOS-model as well. The periodicity relations (2.75) imply that the entire functions $Q_{\pm}(u)$ can always be factorized as

$$Q_{\pm}(u) = e^{i\varphi_{\pm}u/\pi} A_{\pm}(u), \quad A_{\pm}(u) = \prod_{k=1}^N h(u - u_k^{\pm}), \quad (3.15)$$

where $u_1^{\pm}, u_2^{\pm}, \dots, u_N^{\pm}$ are zeroes of $Q_{\pm}(u)$ and the elementary factors

$$h(u) = \vartheta_1(u | q^2) \quad (3.16)$$

have the *double* imaginary period¹³. We will assume that the roots $u_k^{\pm}$ lie in the *fundamental domain*

$$0 < \operatorname{Re} u_k^{\pm} < \pi, \quad -\pi|\tau| < \operatorname{Im} u_k^{\pm} < +\pi|\tau|, \quad k = 1, \dots, N. \quad (3.17)$$

There are two independent sets of the Bethe Ansatz equations (2.4),

$$\frac{f(u_k^{\pm} + \eta)}{f(u_k^{\pm} - \eta)} = -e^{4i\eta\varphi_{\pm}/\pi} \frac{A_{\pm}(u_k^{\pm} + 2\eta)}{A_{\pm}(u_k^{\pm} - 2\eta)}, \quad k = 1, 2, \dots, N \quad (3.18)$$

with all upper and with all lower sings for the zeroes of $Q_+(u)$ and $Q_-(u)$, respectively. The relations (2.75) impose the following constrains on the parameters entering (3.15),

$$\varphi_{\pm} = \pm\varphi + N\pi \pmod{2\pi}, \quad (3.19)$$

$$2\tau\varphi_{\pm} = \mp i\psi - 2 \sum_{k=1}^N u_k^{\pm} + N\pi \pmod{2\pi}. \quad (3.20)$$

Combining these relations one obtains

$$\varphi_+ + \varphi_- = -2\pi m_2, \quad \bar{u}_+ + \bar{u}_- = \pi m_1 + 2\pi\tau m_2, \quad m_1, m_2 \in \mathbb{Z}, \quad (3.21)$$

where

$$\bar{u}_+ = \sum_{k=1}^N u_k^+, \quad \bar{u}_- = \sum_{k=1}^N u_k^-. \quad (3.22)$$

¹³This choice allows a uniform treatment of even and odd N , though for even N the zeroes $Q_{\pm}$ split into $N/2$ equidistant pairs with the separation $\pi\tau$. Actually, we found this redundancy quite useful in controlling the consistency of numerical calculations. Such pairing occurs automatically, without placing any constraints on the positions of zeroes.

Note that for any pair of eigenvalues $Q_{\pm}(u)$ the parameter ψ in (2.75) can be regarded as a function of φ . Indeed, the Bethe Ansatz equations (3.18) only contain the exponents $\varphi_{\pm}$, which up to multiples of π coincide with $\pm\varphi$. Once these equations are solved for $u_k^{\pm}$ the exponent ψ is determined from (3.20).

With the above notations the 8V/SOS-model analog of the formula (3.9) can be written as

$$\begin{aligned} G(u) &= + \frac{e^{-2i\varphi+u/\pi} h'(0)}{h(2\bar{u}_+)} \sum_{k=1}^N \frac{e^{2i\varphi+u_k/\pi} f(u_k^+ + \eta)}{Q_+(u_k^+ + 2\eta) Q'_+(u_k^+)} \frac{h(u + 2\bar{u}_+ - u_k^+)}{h(u - u_k^+)} \\ &= - \frac{e^{-2i\varphi+u/\pi} h'(0)}{h(2\bar{u}_+)} \sum_{k=1}^N \frac{e^{2i\varphi+u_k/\pi} f(u_k^+ - \eta)}{Q_+(u_k^+ - 2\eta) Q'_+(u_k^+)} \frac{h(u + 2\bar{u}_+ - u_k^+)}{h(u - u_k^+)} \end{aligned} \quad (3.23)$$

where

$$G(u) = \frac{1}{2iW(\varphi)} \frac{Q_-(u)}{Q_+(u)} \quad (3.24)$$

and the prime now denotes the derivative with respect to u . This formula is a simple consequence of Bethe Ansatz equations (3.18) and the following elegant identity for the elliptic theta functions¹⁴

$$\prod_{k=1}^n \frac{\vartheta_1(x - y_k)}{\vartheta_1(x - z_k)} = \frac{1}{\vartheta_1(\bar{z} - \bar{y})} \sum_{k=1}^n \left(\prod_{j \neq k} \frac{\vartheta_1(z_k - y_j)}{\vartheta_1(z_k - z_j)} \right) \frac{\vartheta_1(z_k - y_k) \vartheta_1(x - z_k + \bar{z} - \bar{y})}{\vartheta_1(x - z_k)} \quad (3.25)$$

Here x is an independent variable, $y_1, \dots, y_n$ and $z_1, \dots, z_n$ are arbitrary constants,

$$\bar{y} = \sum_{k=1}^n y_k, \quad \bar{z} = \sum_{k=1}^n z_k, \quad (3.26)$$

and for brevity the theta-function $\vartheta_1(x|\mathbf{q})$ is written as $\vartheta_1(x)$. To obtain (3.23) one needs to apply the identity (3.25) to the RHS of the quantum Wronskian relation, written in the form

$$G(u - \eta) - G(u + \eta) = \frac{f(u)}{Q_+(u + \eta) Q_+(u - \eta)}. \quad (3.27)$$

The zero-field variant of (3.23) relates $Q_0(u)$ and $\bar{Q}_0(u)$ defined in (2.46). Let u_k denote the zeroes of $Q_0(u)$,

$$Q_0(u) = \prod_{k=1}^N h(u - u_k^{\pm}), \quad \zeta(u) = \frac{d \log h(u)}{du}. \quad (3.28)$$

From (2.52) one can show that

$$\bar{Q}_0(u) = 2 Q_0(u) \left\{ \frac{i u}{\pi} + \sum_{k=1}^N \zeta(u - u_k) u'_k(0) \right\} + \mathbb{C} Q_0(u) \quad (3.29)$$

where $\mathbb{C}$ is an arbitrary constant and

$$u'_k(0) = \left. \frac{du_k}{d\varphi} \right|_{\varphi=0} = \frac{i f(u_k + \eta) W'(0)}{Q_0(u_k + 2\eta) Q'_0(u_k)}. \quad (3.30)$$

The unknown constant $W'(0)$, defined in (2.53), can be expressed through the zeros u_k from the differential equations (4.12).

¹⁴In the trigonometric limit $\mathbf{q} \rightarrow 0$ this identity reduces to the modified variant of Ex.2 on page 140 of [6], obtained if the constant term therein is “smartly” distributed among the terms in the sum.

3.2 Singular eigenvalues

When η/π is a rational number the spectrum of the transfer matrix of the symmetric 8V-model is degenerate (for sufficiently large N). Solutions of the Bethe Ansatz equations (2.4) for degenerate states are not unique. They contain arbitrary continuous parameters, which determine positions of the complete strings (2.27). There is nothing wrong with this. In particular, this is not an indication that the Bethe Ansatz is “incomplete” [15]. Quite to the contrary, as explained in [13], the appearing continuous parameters are precisely those that are needed to describe the embedding of the corresponding eigenvectors into the eigenspace of the degenerate eigenvalue.

It is not immediately clear, however, how the eigenvalues $Q_{\pm}(u)$, which have no ambiguities at generic values of the parameters η and φ , could acquire continuous degrees of freedom for the degenerate states at special values of η and φ . The explanation is simple: the limiting values of $Q_{\pm}(u)$ are not uniquely defined; they depend on the details of the limiting procedure. Here we will not study this phenomenon in its full generality, but give a particular example. For simplicity we consider the 6V-model in a field, where most calculations can be done analytically. Assume the same notations as in Sect.2.5.3. In particular, recall the definitions of the multiplicative spectral parameter x , the field variable z and the parameter q given in (2.83). For generic z and q the transfer matrix of the 6V-model is completely non-degenerate (the usual degeneracy associated with the reversal of all spins is broken in the presence of the field). Consider the case $N = 4$ and $n = 2$. This sector contains six different eigenvectors. The corresponding polynomials $A_{\pm}(x)$, defined in (2.81), are all of the second degree. Let us parameterize them as

$$A_+(x) = 1 + a_1x + a_2x^2, \quad A_-(x) = 1 + b_1x + b_2x^2. \quad (3.31)$$

Substituting these expressions into the quantum Wronskian relation (3.2) with $\rho_+ = \rho_- = 1$, one gets four equations for the unknown coefficients a_1, a_2, b_1, b_2 ,

$$\begin{aligned} 1 &= a_2b_2, \\ z^2 &= \frac{q^2a_1b_2 + 4q + a_2b_1}{q^2a_2b_1 + 4q + a_1b_2}, \\ z^2 &= \frac{q^4b_2 + q^2(a_1b_1 - 6) + a_2}{q^4a_2 + q^2(a_1b_1 - 6) + b_2}, \\ z^2 &= \frac{q^2b_1 + 4q + a_1}{q^2a_1 + 4q + b_1}. \end{aligned} \quad (3.32)$$

Excluding a_2, b_1, b_2 one obtains a six-degree polynomial equation for a_1 , which factors into two linear and two quadratic equations. One of the latter reads

$$q(1 - q^2z^2)a_1^2 + [4q^2(1 - z^2) + z(1 - q^4)]a_1 + 4q(1 - z)(q^2 + z) = 0, \quad (3.33)$$

while the other is obtained by the substitution $z \rightarrow -z$. Altogether one gets six different solutions for the coefficient a_1 . Once it is found, the remaining coefficients are unambiguously determined by (3.32).

Consider the limit

$$\eta \rightarrow \pi/4, \quad \varphi \rightarrow 0. \quad (3.34)$$

In terms of q and z it corresponds to $q \rightarrow i$, $z \rightarrow 1$. In this limit the two eigenvalues of the transfer matrix, $T^{(1)}(x)$ and $T^{(2)}(x)$, corresponding to two different solutions of (3.33), smoothly approach the same value

$$T(x) = 2x^4 - 12x^2 + 2 = 2(x^2 - b_0^2)(x^2 - b_0^{-2}), \quad b_0 = 1 + \sqrt{2}. \quad (3.35)$$

To calculate $Q_{\pm}(u)$ one also needs to describe the first order deviations of η and φ from their limiting values (3.34). This requires two arbitrary small parameters ϵ_1 and ϵ_2 , or just one such parameter and the ratio ϵ_1/ϵ_2 . We found it convenient to use the following parametrization

$$\eta = \frac{\pi}{4} + (b^4 - 6b^2 + 1)\epsilon + O(\epsilon^2), \quad \varphi = 8(b^4 - 1)\epsilon + O(\epsilon^2) \quad (3.36)$$

where $\epsilon \rightarrow 0$ and b is an arbitrary complex parameter, which is kept finite. The coefficients of the linear terms in ϵ have been specifically chosen to simplify subsequent expressions. The corresponding expansions for q and z simply follow from (2.83). Substituting them into Eq.(3.33), solving it for a_1 and then determining the remaining coefficients in (3.31) from (3.32), one obtains

$$\begin{aligned} A_+^{(1)}(x) &= 1 - \frac{x^2}{b^2} + O(\epsilon), & A_-^{(1)}(x) &= 1 - b^2 x^2 + O(\epsilon), \\ A_+^{(2)}(x) &= 1 - \frac{x^2}{\bar{b}^2} + O(\epsilon), & A_-^{(2)}(x) &= 1 - \bar{b}^2 x^2 + O(\epsilon), \end{aligned} \quad (3.37)$$

where $\bar{b}$ is related to b by a self-reciprocal transformation

$$\bar{b}^2 = \frac{3 - b^2}{1 - 3b^2}, \quad (3.38)$$

which exchanges the two solutions. The corresponding eigenvalues and eigenvectors of the transfer matrix read

$$\begin{aligned} T^{(1)}(x) &= T(x) - 16(1 + b^2)^2 x(1 + x^2)\epsilon + O(\epsilon^2), \\ T^{(2)}(x) &= T(x) + 32(1 - b^2)^2 x(1 + x^2)\epsilon + O(\epsilon^2), \end{aligned} \quad (3.39)$$

where $T(x)$ is given by (3.35) and

$$|\Psi^{(a)}\rangle = |\uparrow\uparrow\uparrow\uparrow\rangle - |\downarrow\uparrow\uparrow\uparrow\rangle + \rho^{(a)}\{|\downarrow\downarrow\uparrow\uparrow\rangle + |\uparrow\uparrow\downarrow\downarrow\rangle - |\uparrow\uparrow\uparrow\downarrow\rangle - |\uparrow\downarrow\downarrow\uparrow\rangle\} + O(\epsilon), \quad a = 1, 2, \quad (3.40)$$

with

$$\rho^{(1)} = \frac{i(1 - b^2)}{1 + b^2}, \quad \rho^{(2)} = \frac{i(1 + b^2)}{2(b^2 - 1)}. \quad (3.41)$$

Here we used self-explanatory notations for the “spin up”, $|\uparrow\rangle$, and “spin down”, $|\downarrow\rangle$, states of the edge spins.

Note, that the case $\eta = \pi/4$ corresponds $m = 1$ and $L = 2$ in (2.20). In terms of the variable x the complete string (2.27) for this case consists of two roots x_1, x_2 , constrained by one relation $x_2 = -x_1$. Obviously, at $\epsilon = 0$ the zeroes of $A_{\pm}(x)$, given by (3.37), are the complete 2-strings at $x_1 = b^{\pm 1}$ or $x_1 = \bar{b}^{\pm 1}$. Their positions are given in terms of the very same parameter b , which defines the embedding of the eigenvectors (3.40) into the eigenspace of the degenerate eigenvalue (3.35) in the sector with two spins up. Remind that the parameter b can be arbitrary; it determines the direction of the 2-variable limit (3.36).

It is interesting to see what happens in some particular cases of (3.36). First, consider the zero-field case when $\varphi \equiv 0$ from the very beginning and η is approaching the value $\pi/4$. In Eq.(3.36) this corresponds $b^2 = \pm 1$. As obvious from (3.37) and (3.38) the limiting values of

$A_{\pm}(x)$ then coincide and the complete 2-string can take one of the two fixed positions with $x_1 = 1$ or $x_1 = i$.

The second interesting case is when η is set to its rational value $\eta \equiv \pi/4$ while the φ is approaching zero from arbitrary values. In Eq.(3.36) this corresponds to $b = b_0^{\pm 1}$, where b_0 is defined in (3.35). Note, also the corresponding value of $\bar{b}_0^2 = b_0^{-2}$. We see that the complete 2-strings in this case occupy precisely the distinguished positions determined by Eq.(2.59). Indeed, taking into account (3.35) it is easy to see that Eq.(2.59) is satisfied with $C(0) = 2$ and $\bar{Q}_{\pm}(x) \equiv A_{\pm}(x)$ (it does not matter which solution in (3.37) is taken, because $A_{\pm}^{(1)}(x) = A_{\mp}^{(2)}(x)$ when $b = b_0$). We would like to stress that the distinguished string positions arising from (2.59) can only be achieved by considering the limit from arbitrary values of the field with fixed rational values of η .

Finally, note that degenerate eigenstates in the 6V- and 8V-models were also considered in [13, 15, 78]

4 Analytic continuation of the eigenvalues

The eigenvalues of the $\mathbf{T}$ - and $\mathbf{Q}$ - matrices of the 8V/SOS-model have very interesting analytic properties with respect to the field parameter φ . In this section we will study these properties by a combination of analytic and numerical techniques. We consider the disordered regime of the 8V-model with real q and η in the range

$$0 < q < 1, \quad \pi/4 < \eta < \pi/2. \quad (4.1)$$

Note that the parameter τ in this case is purely imaginary, $\tau = i|\tau|$.

4.1 Bethe Ansatz equation

We will show that different eigenvalues of the transfer matrix of the symmetric 8V-model can be obtained from each other by the analytic continuation in the variable φ . In doing this we will, obviously, need to consider φ as an independent variable. Therefore, for the eigenvalues $Q_{\pm}(u)$ we assume the SOS-model periodicity (2.75) which allows arbitrary values of φ . In practice it is more convenient to work with a simply related variable, which interpolate the values of φ_{+} (remind that it is related to φ by (3.19)). We will denote this variable by the “upright” symbol ϕ . The Weierstrass factorization for the eigenvalues was discussed in Sect.3.1.2. It is convenient to parameterize the zeroes of $Q_{\pm}(u)$ as

$$u_k^{\pm} = \pi/2 + i\alpha_k^{\pm}. \quad (4.2)$$

Note, that the product representations (3.15) has an obvious ambiguity. For example, the translation of any single root α_k^{+} by the period $2\pi|\tau|$ complemented by the 2π -shift of φ_{+} ,

$$\alpha_k^{+} \rightarrow \alpha_k^{+} - 2\pi i\tau, \quad \varphi_{+} \rightarrow \varphi_{+} + 2\pi, \quad (4.3)$$

leaves $Q_{+}(u)$ unchanged (more precisely, $Q_{+}(u)$ acquires an inessential scalar factor). Using this freedom one can always bring all zeroes of $Q_{\pm}(u)$ to the periodicity rectangular

$$-\pi|\tau| \leq \operatorname{Re} \alpha_k^{\pm} \leq +\pi|\tau|, \quad -\pi/2 \leq \operatorname{Im} \alpha_k^{\pm} \leq \pi/2, \quad k = 1, \dots, N. \quad (4.4)$$

With this convention the values of the exponents $\varphi_{\pm}$ in (3.15) cannot, in general, be restricted to any finite domain.

Introduce two functions

$$\Phi_1(\alpha) = \frac{1}{2\pi i} \log \frac{\vartheta_3(\eta + i\alpha|q)}{\vartheta_3(\eta - i\alpha|q)}, \quad \Phi_2(\alpha) = \frac{1}{2\pi i} \log \frac{\vartheta_1(2\eta + i\alpha|q^2)}{\vartheta_1(2\eta - i\alpha|q^2)}. \quad (4.5)$$

Consider the Bethe Ansatz equations of the following general form

$$N\Phi_1(\alpha_k) - \sum_{j=1}^N \Phi_2(\alpha_k - \alpha_j) = -n_k + \frac{2\eta\phi}{\pi^2}, \quad (4.6)$$

where the numbers $n_1, n_2, \dots, n_N$ and ϕ are given, while the (complex) variables $\alpha_1, \alpha_2, \dots, \alpha_N$ are considered as unknowns. The numbers n_k take half-odd integer values for even N and integer values for odd N . Taking into account (3.15) and (4.2) it is easy to see that the logarithmic form of the Bethe Ansatz equations (3.18) for $Q_{\pm}(u)$ coincide with (4.6) provided α_k and ϕ therein are replaced by $\alpha_k^{\pm}$ and $\varphi_{\pm}$.

Obviously, the numbers $\{n_k\}$ depend on the choice of branches of logarithms in (4.5). In the considered case of the 8V-model, which involves elliptic functions, an extra care is required to define these branches. Since the functions $\Phi_{1,2}$ are periodic

$$\Phi_i(\alpha + i\pi) = \Phi_i(\alpha), \quad i = 1, 2 \quad (4.7)$$

one only needs to specify them in the periodicity strip $0 \leq \text{Im}\alpha < \pi$. We choose the cuts

$$\left\{ \left(m + \frac{1}{2} \right) \pi |\tau| + i \left(\frac{\pi}{2} + n\pi - \eta \right), \left(m + \frac{1}{2} \right) \pi |\tau| + i \left(\frac{\pi}{2} + n\pi + \eta \right) \right\}, \quad m, n \in \mathbb{Z} \quad (4.8)$$

for the function $\Phi_1(\alpha)$ and the cuts

$$\{ m\pi|\tau| + i(n\pi - 2\eta), m\pi|\tau| + i((n-1)\pi + 2\eta) \}, \quad m, n \in \mathbb{Z} \quad (4.9)$$

for $\Phi_2(\alpha)$ (the latter are shown in Fig. 1). We will assume that the values of (4.5) appearing in




Figure 1: The principal Riemann sheet of the function $\Phi_2(\alpha)$.

(4.6) are always taken on the principal sheet of the corresponding Riemann surface defined by the condition

$$\Phi_i(-\alpha) = -\Phi_i(\alpha), \quad i = 1, 2. \quad (4.10)$$

The values on the cuts are taken from the right(left) side for the upper(lower) cuts in Fig.1, as shown by arrows. Note that with these definitions the values of $\Phi_1(\alpha)$ and $\Phi_2(\alpha)$ on the real axis of α are unbounded (this does not happen in the trigonometric case). They monotonically decrease (or increase) with the increase of α . We should warn the reader that this property is never achieved with *ad hoc* computer definitions of the logarithms in (4.5).

Every eigenvalue of the transfer matrix corresponds to some set¹⁵ of the numbers $\{n_k\}$. Even when the branches of the logarithms are completely fixed, there is always an ambiguity in $\{n_k\}$ related to the re-definition of ϕ . Indeed, the transformation

$$\alpha_k \rightarrow \alpha_k, \quad n_k \rightarrow n_k + m, \quad \phi \rightarrow \phi + m \frac{\pi^2}{2\eta}, \quad m \in \mathbb{Z}, \quad (4.11)$$

¹⁵The numerical examples considered below suggest that these sets are the same for $\mathcal{Q}_+(u)$ and $\mathcal{Q}_-(u)$, provided the phases of the logarithms in (4.6) are defined as described above.

leaves Eq.(4.6) unchanged.

Further, the numbers $\{n_k\}$ do not uniquely determine solutions of (4.6). Different solutions corresponding to different eigenvalues could be related with the same set $\{n_k\}$ ¹⁶. In any case once the numbers n_k are fixed the roots $\alpha_k = \alpha_k(\phi)$ solving (4.6) become functions of the complex variable ϕ . It turns out that the latter are multivalued functions with algebraic branching points. Below we will investigate their monodromy properties for various eigenvalues of the transfer matrix.

4.2 Differential form of the Bethe Ansatz equations

The behavior of the roots $\alpha_k(\phi)$ under variations of ϕ can be effectively studied with a differential form of the Bethe Ansatz equations. Differentiating (4.6) with respect to ϕ one immediately obtains a system of ordinary linear differential equations

$$\sum_{k=1}^N A(\alpha)_{jk} \frac{\partial \alpha_k(\phi)}{\partial \phi} = \frac{2\eta}{\pi^2}, \quad j = 1, \dots, N, \quad (4.12)$$

The N by N matrix $A(\alpha)_{jk}$ is given by

$$A(\alpha)_{jk} = \Phi'_2(\alpha_j - \alpha_k) + \delta_{jk} \{ N \Phi'_1(\alpha_j) - \sum_{l=1}^N \Phi'_2(\alpha_j - \alpha_l) \} \quad (4.13)$$

where $\Phi'_{1,2}(\alpha)$ denote derivatives of the functions (4.5) with respect to their argument α .

Using the identity

$$\frac{\theta'_k(x+y|\mathbf{q})}{\theta_k(x+y|\mathbf{q})} + \frac{\theta'_k(x-y|\mathbf{q})}{\theta_k(x-y|\mathbf{q})} = \frac{\theta'_4(2x|\mathbf{q})}{\theta_4(2x|\mathbf{q})} + \sigma_k \theta_2(0|\mathbf{q}) \theta_3(0|\mathbf{q}) \frac{\theta_1(2x|\mathbf{q}) \theta_{5-k}(x+y|\mathbf{q}) \theta_{5-k}(x-y|\mathbf{q})}{\theta_4(2x|\mathbf{q}) \theta_k(x+y|\mathbf{q}) \theta_k(x-y|\mathbf{q})} \quad (4.14)$$

where $k = 1, 2, 3, 4$ and $\sigma_{1,4} = -\sigma_{2,3} = 1$, one can show that $\Phi'_{1,2}(\alpha)$ are meromorphic double-periodic function of the variable α ,

$$\Phi'_1(\alpha) = \xi - \chi \frac{\vartheta_3(0|\mathbf{q}) \vartheta_2(\eta + i\alpha|\mathbf{q}) \vartheta_2(\eta - i\alpha|\mathbf{q})}{\vartheta_2(0|\mathbf{q}) \vartheta_3(\eta + i\alpha|\mathbf{q}) \vartheta_3(\eta - i\alpha|\mathbf{q})}, \quad (4.15)$$

$$\Phi'_2(\alpha) = \xi + \chi \frac{\vartheta_2(2\eta|\mathbf{q}) \vartheta_4(i\alpha|\mathbf{q})}{2\vartheta_1(2\eta + i\alpha|\mathbf{q}^2) \vartheta_1(2\eta - i\alpha|\mathbf{q}^2)}, \quad (4.16)$$

where two constants ξ, χ are given by

$$\xi = \frac{\theta'_4(2\eta|\mathbf{q})}{2\pi \theta_4(2\eta|\mathbf{q})}, \quad \chi = \frac{\theta_2^2(0|\mathbf{q}) \theta_1(2\eta|\mathbf{q})}{2\pi \theta_4(2\eta|\mathbf{q})}. \quad (4.17)$$

For our purposes we need to study the differential equations (4.12) only with the initial conditions corresponding to some eigenvalue of the symmetric 8V-model (typically the ground state eigenvalue). In this case the matrix $A(\alpha)$ is, in general, non-singular. Eq.(4.12) can be then solved for the derivatives $\partial_\phi \alpha_k(\phi)$ and defines locally analytic functions $\alpha_k(\phi)$ of the variable ϕ . Exceptions occur at certain root configurations, corresponding to special values of ϕ , where

¹⁶With the exception of a few simple eigenvalues (e.g. the band of largest eigenvalues [79]) these sets are very poorly understood and their enumeration is a difficult (but, perhaps, not hopeless) problem. To our knowledge the only case where this enumeration problem was completely solved [80] is the $c < 1$ conformal field theory (CFT) with $\Delta \rightarrow \infty$. This CFT can be obtained in the scaling limit of the eight-vertex (six-vertex) model.

the matrix $A(\alpha)$ becomes singular. These singular configurations correspond to the branching points of the solutions in the complex ϕ -plane.

Let us rewrite (4.12) in the form

$$\det \|A(\alpha)\| \frac{\partial \alpha_j(\phi)}{\partial \phi} = \frac{2\eta}{\pi^2} \sum_{k=1}^N A_{jk}^{\text{adj}}(\alpha), \quad (4.18)$$

where A^{adj} stands for the adjoint matrix and assume $\alpha(\phi_0) = (\alpha_1(\phi_0), \dots, \alpha_N(\phi_0))$ is such that

$$\det \|A(\alpha(\phi_0))\| = 0. \quad (4.19)$$

The type of the branching depends on the rank of $A(\alpha(\phi_0))$. The most important case is when

$$\text{rank } A(\alpha(\phi_0)) = N - 1. \quad (4.20)$$

Expanding the determinant $\det \|A(\alpha(\phi))\|$ near the point $\phi = \phi_0$, one obtains

$$\det \|A(\alpha(\phi))\| = \sum_{i=1}^N c_i (\alpha_i(\phi) - \alpha_i(\phi_0)) + \text{higher order terms}, \quad (4.21)$$

where c_i denote some constants. Putting this back into (4.18) and keeping first order terms only, one gets

$$\frac{\partial \alpha_j(\phi)}{\partial \phi} \sum_{i=1}^N c_i (\alpha_i(\phi) - \alpha_i(\phi_0)) = B_j \quad (4.22)$$

where B_j denotes the RHS of (4.18) evaluated at $\phi = \phi_0$. It is easy to see that all B_j are non-zero, otherwise the rank of $A(\phi_0)$ would have been less than $N - 1$. Summing (4.22) over j with the coefficients c_j one obtains

$$\frac{\partial}{\partial \phi} \left(\sum_{i=1}^N c_i (\alpha_i(\phi) - \alpha_i(\phi_0)) \right)^2 = 2 \sum_{i=1}^N c_i B_i. \quad (4.23)$$

Integrating over ϕ and substituting the result back to (4.21) one gets

$$\det \|A(\alpha)\| = \left(2 \sum_{i=1}^N c_i B_i \right)^{1/2} \sqrt{\phi - \phi_0} + O((\phi - \phi_0)) \quad (4.24)$$

Finally, from (4.22) it follows that at $\phi \rightarrow \phi_0$

$$\alpha_j(\phi) - \alpha_j(\phi_0) = D_j \sqrt{\phi - \phi_0} + O((\phi - \phi_0)), \quad D_j = 2B_j \left(2 \sum_{i=1}^N c_i B_i \right)^{-1/2}, \quad (4.25)$$

Therefore, the condition (4.20) implies the square root branching points for all α_j . In principle, higher-order branching points are possible if the rank of the matrix $A(\phi_0)$ drops below $N - 1$. However, we found that the solutions of the Bethe Ansatz equations for $N = 2, 3, 4$ contain the second order branching points only.

Note, that for even N the eigenvalues $Q_{\pm}(u)$ for non-degenerate states possess the more restrictive periodicity (2.72). Their imaginary (quasi-)period is equal to $\pi\tau$ rather than $2\pi\tau$. This is also true for the SOS-model, see Eq.(2.76). Therefore, one can split the roots $\alpha_k(\phi)$ into

the two subsets $\{\alpha_1, \alpha_2, \dots, \alpha_{N/2}\}$ and $\{\alpha_{N/2}, \alpha_{N/2+1}, \dots, \alpha_N\}$ obtained from each other by a uniform shift of all roots by the period $\pi|\tau|$,

$$\alpha_{N/2+k}(\phi) - \alpha_k(\phi) = \pi|\tau|, \quad k = 1, 2, \dots, N/2. \quad (4.26)$$

It is easy to see that under this substitution the system (4.6) reduces to only $N/2$ equations, provided the phases $\{n_{N/2+1}, \dots, n_N\}$ are properly fixed in terms of $\{n_1, \dots, n_{N/2}\}$. Further, the constraints (4.26) are also compatible with the differential equations (4.12). If these constraints hold for the initial data they will continue to hold for an arbitrary ϕ . Therefore for even N one needs to consider only a half of the Bethe roots, for instance, the first $N/2$ roots. The matrix $A(\alpha)$ in (4.12) effectively reduces to a $N/2 \times N/2$ matrix. This considerably simplifies the numeric analysis for even values of N .

4.3 Overview of the procedure

Below we will give a complete classification of the eigenvalues for the symmetric 8V-model with $N = 2, 3, 4$. In most cases we present explicit analytic expressions for the eigenvalues obtained by a direct diagonalization of the transfer matrix. Whenever purely analytic results are not possible, we provide required numeric values for a particular choice of the parameters $\mathbf{q}$ and η . Depending on convenience we will use either of the transfer matrices $\mathbf{T}(u)$ and $\mathbf{T}^B(v)$ related by (2.64), assuming that the parameters u and v are always connected by (2.63). The corresponding eigenvalues, denoted here by $\Lambda(u)$ and $\Lambda^B(v)$ respectively, are related as

$$\Lambda(u) = (-iq^{1/4})^N e^{iuN} \Lambda^B(v), \quad v = u + \frac{\pi\tau}{2}. \quad (4.27)$$

In the previous sections we also considered the eigenvalues of the “higher” transfer matrices $\mathbf{T}_k(u)$, defined by (2.5). In this Section we will denote them as $\Lambda^{(k)}(u)$. With these notations $\Lambda(u) \equiv \Lambda^{(2)}(u)$.

The transfer matrix of the 8V-model commutes [81, 82] with the XYZ-Hamiltonian

$$\mathbf{H}_{XYZ} = -\frac{1}{2} \sum_{j=1}^N (J_x \sigma_1^{(j)} \sigma_1^{(j+1)} + J_y \sigma_2^{(j)} \sigma_2^{(j+1)} + J_z \sigma_3^{(j)} \sigma_3^{(j+1)}) \quad (4.28)$$

provided the constants J_x, J_y, J_z satisfy (A.22). Below we will assume the normalization

$$J_x = 1. \quad (4.29)$$

The eigenvalues of $\mathbf{H}_{XYZ}$ can be found from (A.29) which relates this Hamiltonian to the logarithmic derivative of the transfer matrix $\mathbf{T}^B(v)$.

For each N there are 2^N eigenvalues $\Lambda_i(u)$, $i = 0, 1, \dots, 2^N - 1$. Below we will assume a definite ordering of $\Lambda_i(u)$. We will first split them according to the momenta $\mathcal{P}$ of the corresponding eigenstates¹⁷. Then, in each sector with a fixed value of $\mathcal{P}$ we will arrange $\Lambda_i(u)$ according to eigenvalues of the XYZ-Hamiltonian (4.28).

For every $i = 0, \dots, 2^N - 1$ let $\{\alpha^{(i)}\}$ denote the solution of (4.6), corresponding to the eigenvalue Λ_i . We will say that the contour $C_{ij} \equiv C_{ij}(\zeta)$, $\zeta \in [0, 1]$ connects two eigenvalues Λ_i and Λ_j (corresponding to the values ϕ_i and ϕ_j) if

$$C_{ij}(0) = \phi_i, \quad C_{ij}(1) = \phi_j, \quad \{\alpha^{(i)}\} \xrightarrow{C_{ij}} \{\alpha^{(j)}\}, \quad \Lambda_i \xrightarrow{C_{ij}} \pm \Lambda_j, \quad (4.30)$$

¹⁷The momenta $\mathcal{P}$ are eigenvalues of the *shift operator*, $\mathbf{T}^B(\eta)$ which cyclically shifts the edge spins $\sigma^j \rightarrow \sigma^{j-1}$ to the left by one lattice step.

where we allowed for an ambiguity which can be compensated by an overall shift of all the numbers n_k . Indeed, the only effect of the transformation (4.11) is that the corresponding eigenvalue of the transfer matrix $\Lambda(u)$ acquires an extra sign factor

$$\Lambda(u) \rightarrow (-1)^m \Lambda(u) . \quad (4.31)$$

As a simple illustration consider the ground state eigenvalue. In the regime (4.1) all roots $\{\alpha_k\}$ are real, so they can be arranged in an increasing sequence $\alpha_1 < \alpha_2 < \dots < \alpha_N$. For the ground state the numbers n_k take consecutive integer or half-an-odd integer values

$$n_k = k - \frac{N+1}{2}, \quad k = 1, \dots, N. \quad (4.32)$$

Further, for even N the ground state eigenvalue $\Lambda_0(u)$ of the 8V-model is non-degenerate. It corresponds to $\phi = 0$. At this point the roots solving (4.6) are symmetrically distributed on the interval $(-\pi|\tau|, \pi|\tau|)$, such that $\alpha_k(0) = -\alpha_{N-k+1}(0)$. Let us take this configuration as an initial condition for (4.12) and trace its evolution for real ϕ . When ϕ increases in the positive direction all the roots monotonously move in the negative direction. At $\phi = \pi$ the leftmost root reaches the boundary of the periodicity rectangular (4.4), $\alpha_1(\pi) = -\pi|\tau|$, while one of the middle roots hits the origin $\alpha_{N/2+1}(\pi) = 0$. The remaining $N-2$ roots arrange symmetrically around the origin. The value $\phi = \pi$ belongs to the set of the exponents (2.68) of the symmetric 8V-model. The resulting solution of the Bethe Ansatz equation corresponds to the next-to-leading eigenvalue of the transfer matrix, $\Lambda_1(u)$. Thus, the contour C_{01} in this case is a straight line from $\phi = 0$ to $\phi = \pi$. Further, at $\phi = 2\pi$ the first root becomes

$$\alpha_1(2\pi) = \alpha_N(0) - 2\pi|\tau| , \quad (4.33)$$

while the remaining roots take the original $\phi = 0$ positions, but shifted one step left,

$$\alpha_k(2\pi) = \alpha_{k-1}(0), \quad k = 2, \dots, N . \quad (4.34)$$

The whole process is illustrated in Fig. 2. With an account of (4.3) the resulting root configuration is completely equivalent to initial one at $\phi = 0$. Thus the eigenvalue $\Lambda_0(u)$ is a periodic function of ϕ with the period 2π .




Figure 2: The ϕ -dependence of the Bethe roots for the ground state at $N = 8$.

It is instructive to examine what happens to the exponent ψ in (2.75) for the same course variations of ϕ . It can be calculated from (3.20). We plotted the result in Fig.3. It is a 2π -periodic function which vanishes at the values $\phi = k\pi$, $k \in \mathbb{Z}$, corresponding the symmetric 8V-model, in accordance with (2.68).




Figure 3: The dependence between the exponents ψ on ϕ for the ground state eigenvalue at $N = 8$.

In the remainder of this section we analyze the solutions of the Bethe Ansatz equations for the small size chains with $N \leq 4$. All numerical calculations are performed for the following case

$$q = \frac{1}{10}, \quad \eta = \frac{7\pi}{20}. \quad (4.35)$$

This choice of the parameter η is motivated by the following considerations. First, it allows us to test the behavior of Bethe roots near the special point $\eta = \pi/3$, since the value (4.35) is quite close to it, $\eta - \pi/3 \approx 0.05$. Second, it provides us with all advantages of the rational case without running into the problems of degenerate states. Indeed, the value (4.35) corresponds to $m = 7$ and $L = 10$ in (2.20), while the degenerate states can only occur for $N > L$. Below we will apply the following procedure:

- (i) Analytically diagonalize the transfer matrix of the symmetric 8V-model and thus determine the eigenvalues $\Lambda_i(u)$, $i = 1, \dots, 2^N$.
- (ii) Calculate the corresponding higher eigenvalues $\Lambda_i^{(10)}(u)$ using the formula (2.8).
- (iii) Use the functional relations (2.31) or (2.58) with $L = 10$ to find zeroes of eigenvalues $Q_{\pm}(u)$. For even N this is straightforward since these eigenvalues coincide, however, for odd N one needs to correctly share zeroes between $Q_+(u)$ and $Q_-(u)$.
- (iv) Find the branching points of the eigenvalues with respect to the field variable ϕ .
- (v) On the Riemann surface of the largest eigenvalue $\Lambda_0(u)$ try to find paths connecting it to other eigenvalues $\Lambda_i(u)$ with the exponents from the discrete set (2.68) of the symmetric 8V-model.

The case $N = 2$ is simple and does not really require the steps (ii)-(iii).

The results are presented in tables of two types. In the first one we list eigenvalues of operators $\mathcal{P}$, $\mathcal{S}$, $\mathcal{R}$ and $\mathbf{H}_{XYZ}$ (the latter are denoted as $\mathbf{E}_i$) for each eigenvalue $\Lambda_i(u)$. Remind that we use the normalization (4.29). We also present there numerical values of $\Lambda_i^B = \Lambda_i^B(\pi/2)$ and $\Lambda_i = \Lambda_i(\pi/2)$ (remind that $\Lambda^B(v)$ and $\Lambda(u)$ are related by (4.27)). The second type tables contain the Bethe roots and the values of the phases n_k in (4.6).

Note, that here we did not attempt to diagonalize the infinite dimensional transfer matrices of the SOS-model at arbitrary field and compare the results against the corresponding exact expressions for the eigenvalues. We hope to address this question in the future.

4.4 The case $N = 2$

This case is very simple and all eigenvalues can be calculated analytically. However, it is quite illustrative and reveals some general properties of the Bethe Ansatz equations (4.6).

Four eigenvalues of the transfer-matrix $\mathbf{T}_B(v)$ for $N = 2$ are given by

$$\Lambda_0^B = 2ab + c^2 + d^2, \quad \Lambda_1^B = a^2 + b^2 + 2cd, \quad \Lambda_2^B = a^2 + b^2 - 2cd, \quad \Lambda_3^B = 2ab - c^2 - d^2. \quad (4.36)$$

Corresponding eigenvalues of the Hamiltonian are given by

$$E_0 = -J_x - J_y + J_z, \quad E_1 = -J_x + J_y - J_z, \quad E_2 = J_x - J_y - J_z, \quad E_3 = J_x + J_y + J_z. \quad (4.37)$$

For the symmetric 8V-model with even N the eigenvalues $\mathbf{Q}_i^+(u)$ and $\mathbf{Q}_i^-(u)$ coincide, therefore there is no need to distinguish them. Their analytic expressions read

$$\mathbf{Q}_0(u) = \theta_1(u - \frac{\pi}{2} - \frac{\pi\tau}{2}|q^2)\theta_1(u - \frac{\pi}{2} + \frac{\pi\tau}{2}|q^2) = \frac{\theta_2(0|q)}{2}e^{-i\frac{\pi\tau}{4}}\theta_3(u|q) \quad (4.38)$$

$$\mathbf{Q}_1(u) = e^{iu}\theta_1(u - \frac{\pi}{2} + \pi\tau|q^2)\theta_1(u - \frac{\pi}{2}|q^2) = \frac{\theta_2(0|q)}{2}e^{-i\frac{\pi\tau}{2}}\theta_2(u|q) \quad (4.39)$$

$$\mathbf{Q}_2(u) = e^{iu}\theta_1(u + \pi\tau|q^2)\theta_1(u|q^2) = i\frac{\theta_2(0|q)}{2}e^{-i\frac{\pi\tau}{2}}\theta_1(u|q) \quad (4.40)$$

$$\mathbf{Q}_3(u) = \theta_1(u - \frac{\pi\tau}{2}|q^2)\theta_1(u + \frac{\pi\tau}{2}|q^2) = \frac{\theta_2(0|q)}{2}e^{-i\frac{\pi\tau}{4}}\theta_4(u|q). \quad (4.41)$$

The descriptive properties of the eigenvalues are given Table 1. Table 2 contains the Bethe

i	E_i	Λ_i^B	Λ_i	$\mathcal{P}$	$\mathcal{S}$	$\mathcal{R}$
0	-2.02064769	1.33767166	2.06595956	1	-1	1
1	0.02064769	0.66200143	-0.04265730	1	1	1
2	0.84245604	0.38998228	-0.89156871	1	1	-1
3	1.15754396	-0.28568795	1.21704816	-1	-1	-1

Table 1: Descriptive properties of the eigenvalues for $N = 2$.

roots, the values of the field parameter $\phi = \varphi_+ = \varphi_-$ and the phases n_k for all eigenvalues.

i	α_1	α_2	ϕ	n_1, n_2
0	$i\pi\tau/2$	$-i\pi\tau/2$	0	$\{-1/2, 1/2\}$
1	$i\pi\tau$	0	π	$\{-1/2, 1/2\}$
2	$i\pi\tau - i\pi/2$	$-i\pi/2$	π	$\{-1/2, 1/2\}$
3	$i\pi\tau/2 - i\pi/2$	$-i\pi\tau/2 - i\pi/2$	0	$\{-3/2, -1/2\}$

Table 2: Roots and phases for $N = 2$.

Despite being entire functions of the spectral parameter u , the eigenvalues $\Lambda_i(u)$ and $\mathbf{Q}_i(u)$ are multivalued functions of the field variable ϕ . Actually, for $N = 2$ their analytic properties are relatively simple since they are determined by the properties of only one Bethe root $\alpha_1(\phi)$. Indeed, with an account of (4.26) the Bethe Ansatz equations (4.6) reduce to a single equation

$$2\Phi_1(\alpha_1) = \frac{2\eta}{\pi^2}(\phi + \pi) - (n_1 + \frac{1}{2}), \quad (4.42)$$

which defines the function $\alpha_1(\phi)$. Its branching points are determined by the condition (4.19), which in this case reduces to

$$\Phi'_1(\alpha_1) = 0. \quad (4.43)$$

Solving this equation for α_1 and substituting the result into (4.42) one gets potential locations of the branching points $\phi^{(br)}$ in the ϕ -plane.

$$\phi^{(br)} = \pm\phi_0 + 2\pi m + \frac{\pi^2}{2\eta}n, \quad m, n \in \mathbb{Z} \quad (4.44)$$

where $\Phi'_1(\alpha_1(\phi_0)) = 0$. No explicit analytic expressions for ϕ_0 and $\alpha_1(\phi_0)$ are available. Their asymptotic expansions for small q read

$$\phi_0 = \pi + \frac{2\pi i \sin 2\eta}{\eta} \left[q + q^3 \left(\frac{5}{3} + \frac{\cos 4\eta}{3} \right) + O(q^5) \right], \quad q \rightarrow 0, \quad (4.45)$$

$$\alpha_1(\phi_0) = i\pi\tau - \frac{i\pi}{4} - iq \cos 2\eta + iq^3 \left(\frac{3}{2} \cos 2\eta - \frac{1}{6} \cos 6\eta \right) + O(q^5), \quad q \rightarrow 0. \quad (4.46)$$

The numerical values

$$\phi_0 = \pi + 0.46954959i, \quad \alpha_1(\phi_0) = -0.727625178i + i\pi\tau. \quad (4.47)$$

for the case (4.35) are well approximated by the asymptotic formulae (4.45) and (4.46).

Fig.4 shows two sheets of the Riemann surface of the function $\alpha_1(\phi)$ corresponding to the ground state eigenvalue $\Lambda_0(u)$. The principal sheet is defined by the condition $\alpha_1(0) = i\pi\tau/2$. This sheet contains two cuts $(\phi_0, \pi + i\infty)$ and $(\phi_0^*, \pi - i\infty)$ in the strip $0 < \text{Re } \phi < 2\pi$ (they are shown by solid lines). The same pattern is repeated with the period 2π . The second sheet can be reached from the first one by crossing either of the solid line cuts in Fig.4 from left to right. This sheet contains two new branch cuts (shown by dashed lines) in addition to the same cuts as on the first sheet.

To demonstrate that all the solutions of Bethe Ansatz equations for $N = 2$ can be obtained from each other by an analytic continuation in ϕ we performed numerical calculations using the differential equations (4.12). For $N = 2$ it is a single equation

$$\frac{d\alpha_1}{d\phi} = \frac{\eta}{\pi^2 \Phi'_1(\alpha_1)} \quad (4.48)$$

where $\Phi'_1(\alpha)$ is given by (4.15).

The results are illustrated in Fig.4 by three contours C_{01} , C_{12} and C_{03} , which connect corresponding eigenvalues (we used the definition (4.30)). The contour C_{01} goes between points on the first sheet while the contours C_{12} and C_{03} start on the first and end on the second sheet. Note that the ending point of the contour C_{03} is $\pi^2/(2\eta)$ rather than 0 as it should be according to the Table 2. As explained above, this does not affect the Bethe roots, but changes the sign of the eigenvalue as in (4.31),

$$\Lambda_0 \xrightarrow{C_{03}} -\Lambda_3. \quad (4.49)$$

Apparently it is possible to find a different contour C_{03} ending at the correct point $\phi = 0$ on some other sheet of the same Riemann surface. We leave this as an exercise to the reader.

Note that in the trigonometric limit $q = 0$ the vertical distance between upper and lower cuts vanishes and neighboring cuts join together. In the opposite low temperature limit $q = 1$ all cuts move to infinity and disappear. In both cases the Riemann surface of the eigenvalues splits into disjoint components, though in two different ways. In particular, for $q = 0$ the connected parts of the Riemann surface no longer possess the 2π -periodicity in ϕ .




Figure 4: The principal sheet of the Riemann surface which contains the eigenvalue Λ_0 for $N = 2$.

4.5 The case $N = 3$.

This case is quite important and will be analyzed in more details. When N is odd all eigenvalues of the transfer matrix of the 8V-model are doubly degenerated. So altogether there are four different eigenvalues. Two of them have the momentum $\mathcal{P} = 1$, while for the other two $\mathcal{P} = \omega^{\pm 1}$, $\omega = e^{2\pi i/3}$.

A direct diagonalization of the transfer-matrix leads to the following analytic expressions for the two eigenvalues of $\mathbf{T}_B(v)$ with $\mathcal{P} = 1$,

$$\Lambda_{0,1}^B(v) = \theta_1^3(2\eta|\mathbf{q}) \frac{\theta_1(v|\mathbf{q})\theta_1(v+v_{0,1}|\mathbf{q})\theta_1(v-v_{0,1}|\mathbf{q})}{\theta_1(\eta|\mathbf{q})\theta_1(\eta+v_{0,1}|\mathbf{q})\theta_1(\eta-v_{0,1}|\mathbf{q})} \quad (4.50)$$

where zeros $v_{0,1}$ satisfy the following transcendental equations

$$\frac{\vartheta_4^2(0|\mathbf{q})}{\vartheta_3^2(0|\mathbf{q})} \frac{\vartheta_3^2(v_{0,1}|\mathbf{q})}{\vartheta_4^2(v_{0,1}|\mathbf{q})} = \frac{1 + 7\zeta^2(\gamma - 1) + 3\gamma - 2\zeta \left[1 + \gamma \pm 4\sqrt{\gamma^2 - \zeta\gamma(\gamma + 1) + \zeta^2(1 - \gamma + \gamma^2)} \right]}{(\gamma\zeta - 1)[\zeta(3\gamma + 5) - 3\gamma - 1]}, \quad (4.51)$$

with the constants ζ and γ defined in (A.20). For the choice (4.35) its solutions read¹⁸

$$v_0 = 0.3749001333, \quad v_1 = \frac{\pi}{2} + 0.29695326 i \quad (4.52)$$

The remaining two eigenvalues

$$\Lambda_{2,3}^B(v) = \frac{\theta_1(2\eta|\mathbf{q})\theta_1(v-2\eta|\mathbf{q})\theta_1(v+\eta|\mathbf{q})^2}{\theta_1(\eta|\mathbf{q})} - \omega^{\mp 1} \frac{\theta_1(2\eta|\mathbf{q})^2\theta_1(\eta|\mathbf{q})\theta_1(2v|\mathbf{q})}{\theta_1(v|\mathbf{q})}. \quad (4.53)$$

¹⁸Note that Eq.(4.51) for $\eta = \pi/3$ has a solution $v_0 = 0$, corresponding to the simple ground state eigenvalue $\Lambda_0^B(v) = \theta_1(v|\mathbf{q})^N$, considered in [16, 17].

i	E_i	Λ_i^B	Λ_i	$\mathcal{P}$	$\mathcal{S}$
0	-1.85546785	1.11943180	1.40606193	1	± 1
1	0.69792389	0.25256796	-1.45228936	1	± 1
2	0.57877198	-0.14650972	0.65945338	ω	± 1
3	0.57877198	-0.14650972	0.65945338	ω^{-1}	± 1

Table 3: Descriptive properties of the eigenvalues for $N = 3$.

	Zeros in u			$\phi_{\pm}$	n_1, n_2, n_3
$Q_0^+(u)$	$\frac{\pi}{2} - \frac{\pi\tau}{2}$	$\frac{\pi}{2} + \frac{\pi\tau}{2} - 0.42746i$	$\frac{\pi}{2} + \frac{\pi\tau}{2} + 0.42746i$	$-\frac{\pi}{2}$	$\{-1, 0, 1\}$
$Q_0^-(u)$	$\frac{\pi}{2} - \frac{\pi\tau}{2} - 0.42746i$	$\frac{\pi}{2} - \frac{\pi\tau}{2} + 0.42746i$	$\frac{\pi}{2} + \frac{\pi\tau}{2}$	$+\frac{\pi}{2}$	$\{-1, 0, 1\}$
$\Lambda_0(u)$	$\pm \frac{\pi\tau}{2}$	$\pm \frac{\pi\tau}{2} - 0.374900$	$\pm \frac{\pi\tau}{2} + 0.374900$		
$\Lambda_0^{(3)}(u)$	$\pm \frac{\pi\tau}{2}$	$\pm \frac{\pi\tau}{2} - 1.068576$	$\pm \frac{\pi\tau}{2} + 1.068576$		

Table 4: Zeros of the ground state eigenvalues for $N = 3$.

correspond to $\mathcal{P} = \omega^{\pm 1}$. Their numerical values in Table 3 are real and coincide with each other only due to the choice of the symmetric point $v = \pi/2$ where the second term in (4.53) vanishes. For a generic real v these eigenvalues are complex and non-degenerate.

4.5.1 The eigenvalue Λ_0

Consider the ground state eigenvalue Λ_0 . The numerical calculation with Eq.(2.31) referred to in the step (iii) in Sect.4.3 above require a separation of zeroes between $Q^+(u)$ and $Q^-(u)$. Since the total number of zeroes, $2N = 6$, is small we used a simple trial and error method. Taking an arbitrary subset of 3 zeroes for $Q^+(u)$ and substituting the product (3.15) into the TQ-equation (2.14) one gets a relation which must be fulfilled identically in the variable u at some (yet undermined) value of φ_+ . If the subset of zeroes is chosen incorrectly then no such value exist and the process should be repeated.

The numerical results (with the choice (4.35)) are collected in Table 4 and plotted in Fig.5. Each of the eigenvalues $Q^{\pm}(u)$ has exactly N zeros in the periodicity rectangular

$$0 \leq \text{Re}(u) \leq \pi, \quad -\pi|\tau| \leq \text{Im}(u) \leq \pi|\tau|. \quad (4.54)$$

Any of the eigenvalues $\Lambda^{(k)}(u)$ has $2N$ zeros in the same domain (their imaginary period is $\pi\tau$). The corresponding Bethe roots are

$$\alpha_1^+ = -\frac{1}{2}\pi|\tau|, \quad \alpha_2^+ = \frac{1}{2}\pi|\tau| - q_0, \quad \alpha_3^+ = \frac{1}{2}\pi|\tau| + q_0, \quad q_0 = 0.427465646. \quad (4.55)$$

and the field parameter $\phi_+ = -\pi/2$. The roots α_k^- can be obtained from α_k^+ by either a reflection $\alpha_k^- = -\alpha_{N-k}^+$ or a shift

$$\alpha_k^- = \alpha_{k+1}^+ - \pi|\tau|, \quad k = 1, \dots, N-1, \quad \alpha_N^- = \alpha_1^+ + \pi|\tau|. \quad (4.56)$$

Note also that the sets $\{\alpha_k^+\}$ and $\{\alpha_k^-\}$ precisely interlace each other (see, e.g., Fig.10). The above remarks relating zeroes of the ground state eigenvalues $Q_0^+(u)$ and $Q_0^-(u)$ apply for arbitrary odd N .




Figure 5: Zeros of $Q_0^+(u)$, $Q_0^-(u)$, $\Lambda_0(u)$, $\Lambda_0^{(3)}(u)$ for $N = 3$, marked by $\star$, $\times$, $\bullet$, $\circ$, accordingly.

4.5.2 The eigenvalue Λ_1 .

This is the smallest eigenvalue for $N = 3$ (it has the largest eigenvalue of the Hamiltonian). The numerical zeroes (for the case (4.35)) are presented in Table 5 and plotted in Fig.6. As clearly seen from the picture the zeroes of $Q_1^+(u)$ and $Q_1^-(u)$ form solitary 3-strings.

	Zeros in u			$\phi_{\pm}$	n_1, n_2, n_3
$Q_1^+(u)$	$\frac{\pi}{2} + \frac{\pi\tau}{2} + 0.90837673$	$\frac{\pi}{2} + \frac{\pi\tau}{2}$	$\frac{\pi}{2} + \frac{\pi\tau}{2} - 0.90837673$	$-\frac{3\pi}{2}$	$\{-1, 0, 1\}$
$Q_1^-(u)$	$\frac{\pi}{2} - \frac{\pi\tau}{2} + 0.90837673$	$\frac{\pi}{2} - \frac{\pi\tau}{2}$	$\frac{\pi}{2} - \frac{\pi\tau}{2} - 0.90837673$	$+\frac{3\pi}{2}$	$\{-1, 0, 1\}$
$\Lambda_1(u)$	$\frac{\pi}{2} \pm \frac{\pi\tau}{2} + 0.29655326i$	$\pm \frac{\pi\tau}{2}$	$\frac{\pi}{2} \pm \frac{\pi\tau}{2} - 0.29655326i$		
$\Lambda_1^{(3)}(u)$	$\pm \frac{\pi\tau}{2} + 1.10116874$	$\pm \frac{\pi\tau}{2}$	$\pm \frac{\pi\tau}{2} - 1.10116874$		

Table 5: Zeros for the smallest eigenvalue at $N = 3$.

It is not difficult to check that these solutions of the Bethe Ansatz equations are connected with each other by an analytic continuation in ϕ . Consider the path from $\phi = 3\pi/2$ to $\phi = -3\pi/2$




Figure 6: Zeros of $Q_1^+(u)$, $Q_1^-(u)$, $\Lambda_1(u)$, $\Lambda_1^{(3)}(u)$ for $N = 3$, marked by $\star$, $\times$, $\bullet$, $\circ$, accordingly.

along the real axis of ϕ (with a small imaginary part for a better convergence). It turns out that the solution $\{\alpha_1^-, \alpha_2^-, \alpha_3^-\}$ continued along this path transforms into another solution

$$\{\tilde{\alpha}_k^+\} = \{\alpha_1^+ - 4\pi|\tau|, \alpha_2^+ + 2\pi|\tau|, \alpha_3^+ + 2\pi|\tau|\}, \quad (4.57)$$

where α_i^+ , $i = 1, 2, 3$ correspond to zeros of $Q_1^+(u)$ from the Table 5. Remind that α_k and u_k are related by (4.2). It is easy to check that shifts by $2\pi k|\tau|$ in (4.57) can be compensated by a proper choice of n_k without changing the field $\phi_+ = -3\pi/2$. Table 5 contains the resulting values of n_k corresponding to the roots $\{\alpha_1^+, \alpha_2^+, \alpha_3^+\}$.

The problem of the analytic continuation between different eigenvalues of the transfer matrix is much more complicated. The most difficult part of this procedure is a numerical study of the Riemann surface of the eigenvalues. For $N \geq 3$ the difficulty lies in the fact that the branching condition (4.20) just defines an $(N - 1)$ -dimensional hyper-surface in the space of Bethe roots which, by itself, does not correspond to any particular value of the field ϕ . To obtain the structure of the branching points in the ϕ -plane one needs to determine how this hyper-surface intersects with required solutions of the Bethe Ansatz equations. In practice we used the differential equations (4.12) and numerically studied monodromy properties of the solutions when ϕ was varied around random loops in the complex plane.

Remarkably, we found that the largest and smallest eigenvalues, $\Lambda_0(u)$ and $\Lambda_1(u)$, do indeed correspond to different branches of the same (multivalued) function of ϕ . The contour C_{01} in the complex ϕ -plane connecting these eigenvalues is shown in Fig.7. As evident from the figure, the structure of branching cuts is extremely complicated. This figure shows the (upper half-plane) cuts on just two sheets of the corresponding Riemann surface. The first sheet (solid line cuts) contains the eigenvalue $Q_0^-(u)$ at $\phi = \pi/2$, while the second one (dashed line cuts) contains the

eigenvalue $Q_1^-(u)$ at $\phi = 3\pi/2$. Note that the contour C_{01} leaves the first sheet when it goes under the cut at $\text{Re } \phi = 0$ and arrives to the second of these sheets from under the (dashed line) cut at $\text{Re } \phi = \pi - \alpha$. In between these two points the contour crosses additional cuts on other sheets of the Riemann surface, which are not shown in the picture.




Figure 7: Structure of the cuts on the principal sheet Riemann surface containing Λ_0 (solid lines) and on the sheet containing Λ_1 (dashed lines). Only the upper half-plane is shown as the arrangement of the cuts is reflection-symmetric with respect to the real axis of ϕ

Note that in the limit $\eta \rightarrow \pi/3$ the zeroes of $Q_1^\pm(u)$ form complete 3-strings which cancel out from the TQ-relation. As a result one gets

$$\Lambda_1(u) = -\theta_4^3(u + \frac{\pi}{3}|\mathbf{q}) - \theta_4^3(u - \frac{\pi}{3}|\mathbf{q}), \quad \eta = \frac{\pi}{3}. \quad (4.58)$$

The minus signs arise from the phase factors $\exp(\pm 2i\eta\phi/\pi) = -1$ at $\phi = 3\pi/2$. When $\eta \rightarrow \pi/3$ the parameter α tends to zero (this parameter determines positions of several cuts in Fig.7). In addition to that the dashed line cut with $\text{Re } \phi = \pi$ and its symmetric reflection in the lower half plane both approach the real axes. Therefore, the contour C_{01} gets pinched twice in both vertical and horizontal directions. As a result the complete string solution corresponding to $\Lambda_1(u)$ for $\eta = \pi/3$ is no longer analytically connected to the ground state eigenvalue $\Lambda_0(u)$.

4.5.3 The eigenvalues $\Lambda_{2,3}$

It is enough to consider only one eigenvalue Λ_2 with $\mathcal{P} = \omega$, since all results for Λ_3 can be obtained by a complex conjugation. The analytic formula for $\Lambda_2^B(v)$ is given in (4.53). The Bethe roots and phases for $Q_2^-(u)$ read

$$\{\alpha_1^-, \alpha_2^-, \alpha_3^-\} = \{-0.24023993, -0.45552631 \pm 0.75825341 i\}, \quad \{n_1, n_2, n_3\} = \{0, 0, 1\} \quad (4.59)$$

and the exponent $\varphi_- = \pi/2$. Similarly, for $Q_2^+(u)$ one has

$$\alpha_k^+ = \alpha_k^+ + \pi|\tau|, \quad \varphi_+ = -\frac{5\pi}{2}, \quad \{n_1, n_2, n_3\} = \{0, 0, 1\}. \quad (4.60)$$

For the eigenvalue Λ_3 all $\alpha_i^\pm$, $\varphi_\pm$ and n_k change their sign with respect to those for Λ_2 .

The problem of the analytic continuation for these eigenvalues and, in particular, the question about their connection to the ground state eigenvalue $\Lambda_0(u)$ have not been considered.

4.6 The case $N = 4$.

This is the last case where we systematically study all eigenvalues of the transfer matrix. As before the eigenvalues will be first grouped according to their momenta, which in this case takes four possible values $\mathcal{P} = 1, -1, \pm i$. In total there are 16 eigenvalues, all non-degenerate. Below we will show that all the eigenvalues with $\mathcal{P} = \pm 1$ (there are ten such eigenvalues) correspond to different branches of the same multivalued function of ϕ and explicitly present all paths on the Riemann surface, which connect these eigenvalues to each other.

4.6.1 The sector $\mathcal{P} = 1$.

This sector contains 6 eigenvalues listed in Table 6. Three of them are given by the formula

$$\Lambda_i^B = a^4 + b^4 + 2ab(a^2 + b^2) \left[\frac{x_i + (\gamma + 1)(\zeta - 1)}{1 - \gamma} \right] - 2a^2b^2 \left[\frac{(\gamma + 1)[\zeta^2(\gamma - 1) - (\zeta + 1)x_i] - x_i^2}{(\gamma - 1)^2} \right], \quad (4.61)$$

where $i = 0, 3, 5$ and the constants x_0, x_3 and x_5 satisfy the same cubic equation

$$x^3 + x[2\zeta(\gamma + 1)^2 - (\zeta^2 + 1)(3 - 2\gamma + 3\gamma^2)] + 2(\zeta - 1)^2(\zeta + 1)(\gamma - 1)^2(\gamma + 1) = 0. \quad (4.62)$$

with ζ, γ defined in (A.20). With the choice (4.35) this equation has three real roots

$$x_0 = 12.98056849, \quad x_3 = -3.41421744, \quad x_5 = -9.56635105. \quad (4.63)$$

The remaining three eigenvalues are polynomially expressed through the Boltzmann weights (A.13)

$$\Lambda_1^B = (a^2 + b^2)(c^2 + d^2) + (a^2 + b^2 + c^2 + d^2)(ab + cd) + 4abcd, \quad (4.64)$$

$$\Lambda_2^B = (a^2 + b^2)(c^2 + d^2) + (a^2 + b^2 + c^2 + d^2)(ab - cd) - 4abcd, \quad (4.65)$$

$$\Lambda_4^B = a^4 + b^4 - 2c^2d^2. \quad (4.66)$$

The corresponding eigenvalues of the XYZ-Hamiltonian (4.28) read

$$E_1 = -2J_1, \quad E_2 = -2J_2, \quad E_4 = -2J_3. \quad (4.67)$$

i	E_i	Λ_i^B	Λ_i	$\mathcal{P}$	$\mathcal{S}$	$\mathcal{R}$
0	-2.75822456	1.30674491	2.36255547	1	1	1
1	-2.00000000	0.94124717	0.45441276	1	-1	1
2	-1.17819166	0.61623116	-0.92097245	1	-1	-1
3	0.72548274	0.14757534	-1.33882675	1	1	1
4	0.86310373	0.12908441	-1.21915423	1	1	-1
5	2.03274182	0.05569128	0.61370849	1	1	1

Table 6: Properties of the eigenvalues with $\mathcal{P} = 1$ for $N = 4$.

Table 7 contains positions of the Bethe roots $\alpha_{1,2}$, the values of the field ϕ and phases n_k appearing in (4.6). Note that, since N is even the roots satisfy (4.26) and it is enough to present only two roots $\alpha_{1,2}$.

i	α_1	α_2	ϕ	n_1, n_2, n_3, n_4
0	-1.41965608	-0.88292901	0	$\{-3/2, -1/2, 1/2, 3/2\}$
1	$i\pi\tau$	$i\pi\tau/2$	π	$\{-3/2, -1/2, 1/2, 3/2\}$
2	$i\pi\tau - i\pi/2$	$i\pi\tau/2$	π	$\{-3/2, -1/2, 1/2, 3/2\}$
3	$i\pi\tau - 0.59503973i$	$0.59503973i$	0	$\{-3/2, -1/2, 1/2, 3/2\}$
4	$i\pi\tau - i\pi/2$	0	0	$\{-3/2, -1/2, 1/2, 3/2\}$
5	$-1.65004854 - i\pi/2$	$-0.65253655 - i\pi/2$	0	$\{-5/2, 1/2, -1/2, 5/2\}$

Table 7: Bethe roots, values of the field and phases for $\mathcal{P} = 1$ and $N = 4$.

4.6.2 The sector $\mathcal{P} = -1$.

This sector contains 4 eigenvalues. All of them are polynomial in Boltzmann weights and the corresponding Bethe roots can be found explicitly.

$$\Lambda_6^B = 2a^2b^2 - c^4 - d^4 \quad (4.68)$$

$$\Lambda_7^B = 2(a^2b^2 + c^2d^2) - (a^2 + b^2)(c^2 + d^2) \quad (4.69)$$

$$\Lambda_8^B = -(a^2 + b^2)(c^2 + d^2) + (a^2 + b^2 + c^2 + d^2)(ab - cd) + 4abcd \quad (4.70)$$

$$\Lambda_9^B = -(a^2 + b^2)(c^2 + d^2) + (a^2 + b^2 + c^2 + d^2)(ab + cd) - 4abcd \quad (4.71)$$

The associated eigenvalues of the XYZ-Hamiltonian are

$$E_6 = 2J_3, \quad E_7 = 0, \quad E_8 = 2J_2, \quad E_9 = 2J_1. \quad (4.72)$$

i	E_i	Λ_i^B	Λ_i	$\mathcal{P}$	$\mathcal{S}$	$\mathcal{R}$
6	-0.86310373	-0.51124109	1.25718614	-1	1	-1
7	0	-0.27935395	1.63561756	-1	1	1
8	1.17819166	-0.09456292	0.86905645	-1	-1	-1
9	2.00000000	-0.05570662	-0.54254103	-1	-1	1

Table 8: Properties of eigenvalues for $\mathcal{P} = -1$ and $N = 4$.

i	α_1	α_2	ϕ	n_1, n_2, n_3, n_4
6	$i\pi\tau/2 - i\pi/2$	$i\pi\tau/2$	0	$\{-5/2, -3/2, -1/2, 1/2\}$
7	$i\pi\tau/2 - i(\pi/2 - \eta)$	$i\pi\tau/2 + i(\pi/2 - \eta)$	0	$\{-3/2, -1/2, 1/2, 3/2\}$
8	$i\pi\tau/2 - i\pi/2$	0	$-\pi$	$\{-7/2, -1/2, -3/2, 3/2\}$
9	$i\pi\tau/2 - i\pi/2$	$-i\pi/2$	$-\pi$	$\{-5/2, 1/2, -1/2, 5/2\}$

Table 9: Bethe roots, values of the field and phases with $\mathcal{P} = -1$ at $N = 4$.

Using the differential equations (4.12) we have found that all solutions of the Bethe Ansatz equations (4.6) corresponding to $\mathcal{P} = \pm 1$ can be obtained from the ground state solution by the analytical continuation in ϕ . For the numerical calculations we always assumed the values (4.35). Consider the principal sheet of the Riemann surface containing the ground state solution at $\phi = 0$. As explained before the cut structure on this sheet is always symmetric with respect to

the real axis and periodic with the period 2π . Therefore we consider only the upper half of the periodicity strip. There is only one cut $(\pi + 0.67535i, \pi + i\infty)$ on this sheet shown by solid lines in Fig.8. Encircling the branching point (as in the contour C_{12}) brings one to the second sheet containing four additional cuts (shown by dashed lines in Fig.8). In this figure we also shown the contours which connect Λ_0 with five other eigenvalues $\Lambda_1, \Lambda_2, \Lambda_3, \Lambda_4, -\Lambda_6$. The change of sign for Λ_6 is related with the transformation (4.11) and (4.31) with $m = 1$. The horizontal coordinates of the cuts shown on the figures have been numerically fitted by considering a few different values of η in the vicinity $\eta = 7\pi/20$.




Figure 8: The principal sheet of the Riemann surface containing the eigenvalue Λ_0 for $N = 4$.




Figure 9: The sheet of the Riemann surface containing Λ_4 for $N = 4$.

Paths to the other four eigenvalues $\Lambda_5, \Lambda_7, \Lambda_8$ and Λ_9 are more conveniently described on a different sheet of the Riemann surface which contains the eigenvalue Λ_4 at $\phi = 0$. The structure of cuts on this sheet and connecting contours are shown in Fig.9. Again at the points $\pi^2/(2\eta) - \pi$

and $2\pi - \pi^2/(2\eta)$ we have to take into account (4.31) with $m = \pm 1$. For Λ_7 we used the shift (4.3) to move the Bethe roots to the region (4.4).

4.6.3 The sector $\mathcal{P} = i$.

There are three complex eigenvalues with $\mathcal{P} = i$ and three complex conjugated eigenvalues with $\mathcal{P} = -i$. Again it is sufficient to consider only $\mathcal{P} = i$.

$$\Lambda_{10}^B = 2(a^2b^2 - c^2d^2) + i(a^2 - b^2)(c^2 - d^2) \quad (4.73)$$

$$\Lambda_{11}^B = (ab + cd)(a^2 + b^2 - c^2 - d^2) + i(a^2 - b^2)(c^2 - d^2) \quad (4.74)$$

$$\Lambda_{12}^B = (ab - cd)(a^2 + b^2 - c^2 - d^2) + i(a^2 - b^2)(c^2 - d^2). \quad (4.75)$$

Note that for all of them the corresponding energies E_i of the XYZ-Hamiltonian are zero. In general, these eigenvalues are complex. However, at the symmetric point $v = \pi/2$ where $a = b$, all eigenvalues are real. All information about these eigenvalues is collected in Tables 10, 11. The parameters of the eigenvalues with the momentum $\mathcal{P} = -i$ are obtained by negating of all α_i 's, ϕ 's and n_k 's.

i	E_i	Λ_i^B	Λ_i	$\mathcal{P}$	$\mathcal{S}$	$\mathcal{R}$
10	0	0.12908441	1.25718614	i	1	-1
11	0	-0.09456292	-0.92097245	i	-1	-1
12	0	-0.05570662	-0.54254103	i	-1	1

Table 10: Eigenvalues with $\mathcal{P} = i$ at $N = 4$.

i	α_1	α_2	ϕ	n_1, n_2, n_3, n_4
10	$-1.50555656 - i\pi/2$	-0.79702853	0	$\{-5/2, -1/2, -1/2, 3/2\}$
11	$-1.89851386 - i\pi/2$	-1.55536378	π	$\{-3/2, -3/2, 1/2, 1/2\}$
12	$3i\pi\tau/4 - 0.40566532i$	$3i\pi\tau/4 + 0.40566532i$	π	$\{-3/2, -3/2, 1/2, 1/2\}$

Table 11: Bethe roots, values of the field and phases for $\mathcal{P} = i$ and $N = 4$.

The problem of analytic continuation for these eigenvalues has not been considered.

4.7 The case $N = 13$

The purpose of this section is to analyze a distribution of zeroes for the ground state eigenvalues for sufficiently large value of N . We deliberately choose an odd value of N when $Q_+(u)$ and $Q_-(u)$ are linearly independent and their quantum Wronskian is nonzero. In this section it is more convenient to work with the original Baxter's normalization of eigenvalues $Q_{\pm}^B(v)$ and the variable v

$$Q_{\pm}(u) = e^{ivN/2} Q_{\pm}^B(v), \quad v = u + \pi\tau/2. \quad (4.76)$$

Introduce the linear combinations $Q_{1,2}(v)$,

$$Q_{\pm}^B(v) = (Q_1(v) \pm Q_2(v))/2, \quad (4.77)$$

such that

$$Q_{1,2}(v + \pi) = (-1)^{(N-1)/2} Q_{2,1}(v) \quad (4.78)$$

Numerical zeroes of eigenvalues can be easily calculated from the Bethe Ansatz equations (4.6) with the set of phases (4.32). Here we demonstrate an alternative numerical method which works well for the considered case. For any odd N the ground state eigenvalue $\Lambda_0^B(v)$ an odd function of v , having N zeroes in its periodicity rectangular (2.65). Using the identity

$$2\vartheta_1(x+y|\mathbf{q})\vartheta_1(x-y|\mathbf{q}) = \bar{\vartheta}_4(2x)\bar{\vartheta}_3(2y) - \bar{\vartheta}_3(2x)\bar{\vartheta}_4(2y), \quad \bar{\vartheta}_i(x) \equiv \vartheta_i\left(\frac{x}{2}|\mathbf{q}^{1/2}\right), \quad (4.79)$$

it can be represented as

$$\Lambda_0^B(v) = \vartheta_1(v|\mathbf{q}) \sum_{k=0}^{(N-1)/2} t_k \bar{\vartheta}_3(2v)^k \bar{\vartheta}_4(2v)^{(N-1-2k)/2}, \quad (4.80)$$

where t_k are constants. Similarly, $Q_1(v)$ can be written as

$$Q_1(v) = 2^{-N} \bar{\vartheta}_3(v) \sum_{k=0}^{(N-1)/2} c_k \bar{\vartheta}_3(v)^{2k} \bar{\vartheta}_4(v)^{N-1-2k}, \quad (4.81)$$

with some unknown constants c_k . In [16] we used the representation (4.81) for $\eta = \pi/3$, however, it is valid for the ground state eigenvalues with arbitrary values of η (the case $\eta = \pi/3$ is special because all coefficients c_k can be calculated explicitly [16]). Substituting (4.80) and (4.81) into the TQ-equation and evaluating it numerically for several values of the spectral parameter one gets a bi-linear system of equations for the unknown coefficients t_k and c_k . For the choice (4.35) its numerical solution is given in Table 12.

k	c_k	k	c_k
0	1.00000000	3	-26.42365231
1	-5.63412047	4	28.59503534
2	15.52969161	5	-18.21779332
		6	5.231349086

Table 12: Numerical values of the coefficients c_k in (4.81) for $N = 13$ in the case (4.35).

i	α_i^-	i	α_i^-	i	α_i^-
1	-1.97251218	5	-1.00040409	9	0.91284796
2	-1.50058366	6	-0.80200143	10	1.07830969
3	-1.30218100	7	-0.33007292	11	1.22427541
4	-1.15129255	8	0.64848502	12	1.38973713
				13	1.65410007

Table 13: Bethe roots for the ground state at $N = 13$.

Using these results and (4.76–4.78) one can find the Bethe roots $\alpha_k^\pm$, $k = 1, \dots, 13$. The results for α_k^- are given in Table 13, moreover, $\alpha_i^+ = -\alpha_{N-i+1}^-$ and $\varphi_\pm = \mp\pi/2$.

Further, zeros for $\Lambda_0(u)$ and $\Lambda_0^{(3)}(u)$ are as follows. The $2N = 26$ zeros of $\Lambda_0(u)$ read

$$u_k = \pm \frac{\pi\tau}{2}, \quad \pm \frac{\pi\tau}{2} \pm \{r_1, r_2, r_3, r_1^*, r_2^*, r_3^*\}, \quad (4.82)$$

where the signs can be chosen independently and r_i^* stand for complex conjugates of r_i . Similarly, for $\Lambda_0^{(3)}(u)$,

$$u_k = \pm \frac{\pi\tau}{2}, \quad \pm \frac{\pi\tau}{2} + \frac{\pi}{2} \pm \{s_1, s_2\}, \quad \pm \frac{\pi\tau}{2} + \frac{\pi}{2} \pm \{s_3, s_4, s_3^*, s_4^*\}. \quad (4.83)$$

The numerical constants r_i , s_i are given in Table 14. The zeroes of eigenvalues are plotted in Fig. 10.

k	s_k	r_k
1	0.27442683	$0.40999133 + 0.42208534 i$
2	0.47123771	$0.49344215 + 0.21629447 i$
3	$0.28395326 + 0.15112367 i$	$0.51703097 + 0.067457998 i$
4	$0.32747706 + 0.34260021 i$	

Table 14: Parameters s_k , r_k which determine positions of zeros $\Lambda_0(u)$ and $\Lambda_0^{(3)}(u)$.




Figure 10: Zeros of $Q_0^+(u)$, $Q_0^-(u)$, $\Lambda_0(u)$, $\Lambda_0^{(3)}(u)$ for $N = 13$, marked by $\star$, $\times$, $\bullet$, $\circ$, accordingly.

Note that the sets $\{\alpha_k^+\}$ and $\{\alpha_k^-\}$ precisely interlace each other. This fact looks quite remarkable given these sets are obtained from each other by the uniform translation (4.56) over the distance $\pi|\tau|$ which is much larger than the spacing between roots. For a numerical study of the Bethe Ansatz equations on a large chain of even length (up to $N = 512$) we refer the reader to [83].

5 Conclusion

In this paper we developed some new ideas in the classical subject of Baxter's celebrated eight-vertex and solid-on-solid models. Our primary observation concerns a (previously unnoticed) arbitrary field parameter in the solvable solid-on-solid model. This parameter is analogous to the horizontal field in the six-vertex model. This fact might not be so surprising to experts, since all the hard work has been done before and one just needs to lay side-by-side the papers [4, 11, 12] to realize that an arbitrary field parameter is, in fact, required to describe the continuous spectrum of the unrestricted solid-on-solid model.

The introduction of an arbitrary field allowed us to develop a completely analytic theory of the functional relations in the 8V/SOS-model. The solutions of the Bethe Ansatz equations are multivalued functions of the field variable, having algebraic branching points. It is plausible that many (if not all) eigenvalues of the transfer matrix can be obtained from each other via analytic continuation in this variable. To demonstrate this we performed a comprehensive study of all eigenvalues for the 8V-model for small chains of the length $N \leq 4$ with a combination of analytic and numeric techniques. In particular, we saw in these cases that the largest and smallest eigenvalues of the transfer matrix are always connected by the analytic continuation. This study was partially motivated by our attempts to understand properties of the eigenvalues in the ferromagnetic regime [84], analytically connecting it with the disordered regime. Note that the ferromagnetic regime eigenvalues are important in connection with the AdS/CFT correspondence (see [85] and references therein); we hope to study them elsewhere.

The field parameter is also important in our future considerations [18] of the quantum field theory limit of the 8V/SOS-model, where it becomes the massive sine-Gordon model. Note, that the connection between the largest and next-to-largest eigenvalues in this model was previously studied [53] via the thermodynamic Bethe Ansatz. The authors of [53] also considered the analytic continuation but in a different variable, namely, the scaling variable, which has no a direct analogue in the lattice theory.

It appears that the analytic structure of eigenvalues in the eight-vertex/SOS model certainly deserves further studies. Somewhat simpler (but still very interesting) structure arises in the six-vertex model and, especially, in the $c < 1$ conformal field theory [80]. In the latter case the Riemann surface of the eigenvalues closes within each level subspace of the Virasoro module.

Acknowledgments

The authors thank R.J.Baxter, M.T.Batchelor, B.M.McCoy, K.Fabrizius, P.A.Pearce, S.M.Sergeev, F.A.Smirnov and M.Bortz for useful remarks. One of us (VVB) thanks S.M.Lukyanov and A.B.Zamolodchikov for numerous discussions of the analytic structure of eigenvalues in solvable models. This work was supported by the Australian Research Council.

Appendix A. The eight-vertex model

In this Appendix we briefly summarize basic properties of the symmetric eight-vertex (8V) model used in this paper. For a more detailed information the reader should consult with Baxter's original publications [1–5].

Consider a square lattice of N columns and M rows, with the toroidal boundary conditions. Each edge of the lattice carries a spin variable α taking two values $\alpha = +$ and $\alpha = -$, corresponding to the “spin-up” and “spin-down” states of the edge. Each vertex is assigned with a Boltzmann weight $R(\alpha, \alpha' | \beta, \beta')$ depending on the spin states $\alpha, \alpha', \beta, \beta'$ of the surrounding edges arranged as in Fig.11. There are only eight “allowed” vertex configurations, shown in




Figure 11: The arrangement of four spins around the vertex.

Fig.12, which have non-vanishing Boltzmann weights. These weights are not arbitrary; they are parameterized by only four arbitrary constants a, b, c, d ,

$$\phi_1 = \phi_2 = a, \quad \phi_3 = \phi_4 = b, \quad \phi_5 = \phi_6 = c, \quad \phi_7 = \phi_8 = d. \quad (\text{A.1})$$

The remaining eight configurations are forbidden; their Boltzmann weight is zero.

The partition function

$$Z = \sum_{(\text{spins})} \prod_{(\text{vertices})} R(\alpha, \alpha' | \beta, \beta') \quad (\text{A.2})$$

is defined as the sum over all spin configurations of the whole lattice, where each configuration is counted with the weight equal to the product of vertex weights over all lattice vertices.

The vertex weight matrix $R(\alpha, \alpha' | \beta, \beta')$ can be thought as a matrix acting in the direct product of the two two-dimensional vector spaces $\mathbb{C}^2 \otimes \mathbb{C}^2$, where the indices α, α' refer to the first space and the indices β, β' to the second. It has an elegant representation [1]

$$\mathbf{R} = \sum_{k=1}^4 w_k (\sigma_k \otimes \sigma_k), \quad (\text{A.3})$$

where σ_k , $k = 1, 2, 3, 4$, are the Pauli matrices

$$\sigma_1 = \begin{pmatrix} & 1 \\ 1 & \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} & -i \\ i & \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & \\ & -1 \end{pmatrix}, \quad \sigma_4 = \begin{pmatrix} 1 & \\ & 1 \end{pmatrix}, \quad (\text{A.4})$$




Figure 12: Eight allowed vertex configuration and their Boltzmann weights. Thin lines represent the “spin-up” states and the bold lines represent the “spin-down” states of the edge spins

and

$$w_1 = \frac{c+d}{2}, \quad w_2 = \frac{c-d}{2}, \quad w_3 = \frac{a-b}{2}, \quad w_4 = \frac{a+b}{2}. \quad (\text{A.5})$$

The matrix $\mathbf{R}$ can conveniently be presented as a two-by-two block matrix acting in the first space with the two-dimensional matrix blocks acting in the second space,

$$\mathbf{R} = \begin{pmatrix} \mathbf{R}_{++} & \mathbf{R}_{+-} \\ \mathbf{R}_{-+} & \mathbf{R}_{--} \end{pmatrix} \quad (\text{A.6})$$

where

$$\mathbf{R}_{++} = \begin{pmatrix} a & \\ & b \end{pmatrix}, \quad \mathbf{R}_{+-} = \begin{pmatrix} & d \\ c & \end{pmatrix}, \quad \mathbf{R}_{-+} = \begin{pmatrix} & c \\ d & \end{pmatrix}, \quad \mathbf{R}_{--} = \begin{pmatrix} b & \\ & a \end{pmatrix}. \quad (\text{A.7})$$

The matrix $\mathbf{R}$ possesses simple symmetry relations, obvious from (A.3),

$$\mathbf{R} = (\sigma_k \otimes \sigma_k) \mathbf{R} (\sigma_k \otimes \sigma_k), \quad k = 1, 2, 3. \quad (\text{A.8})$$

The row-to-row transfer matrix $\mathbf{T}$ acts in space of states of N spins located on a horizontal row of vertical edges. It is defined as the trace of the product of the two-by-two matrices (A.6)

$$\mathbf{T} = \text{Tr}_{\mathbb{C}^2} \left(\mathbf{R}^{(1)} \mathbf{R}^{(2)} \dots \mathbf{R}^{(N)} \right), \quad (\text{A.9})$$

where the operator entries (A.7) of the matrix $\mathbf{R}^{(j)}$ act on the states of the j -th spin in the row. The partition function (A.2) can written as

$$Z = \text{Tr} (\mathbf{T}^M), \quad (\text{A.10})$$

where M is the number of rows of the lattice. It follows from (A.8) that

$$[\mathbf{T}, \mathcal{S}] = [\mathbf{T}, \mathcal{R}] = 0 \quad (\text{A.11})$$

where the operators

$$\mathcal{S} = \sigma_3^{(1)} \otimes \sigma_3^{(2)} \otimes \cdots \otimes \sigma_3^{(N)}, \quad \mathcal{R} = \sigma_1^{(1)} \otimes \sigma_1^{(2)} \otimes \cdots \otimes \sigma_1^{(N)} \quad (\text{A.12})$$

defined as the product of the local spin operators $\sigma_3^{(j)}$ and $\sigma_1^{(j)}$ over the all sites, $j = 1, \dots, N$, in a row. Note, that for an odd N these operators do not commute among themselves (indeed, $\mathcal{R}\mathcal{S} = (-1)^N \mathcal{S}\mathcal{R}$) and only one of them can be diagonalized simultaneously with $\mathbf{T}(v)$. Below we will always assume the basis where the operator $\mathcal{S}$ is diagonal (then for even N the operator $\mathcal{R}$ will be diagonal as well).

Following [1] we parameterize the Boltzmann weights a, b, c, d as

$$\begin{aligned} a &= \rho \vartheta_4(2\eta | \mathbf{q}^2) \vartheta_4(v - \eta | \mathbf{q}^2) \vartheta_1(v + \eta | \mathbf{q}^2), \\ b &= \rho \vartheta_4(2\eta | \mathbf{q}^2) \vartheta_1(v - \eta | \mathbf{q}^2) \vartheta_4(v + \eta | \mathbf{q}^2), \\ c &= \rho \vartheta_1(2\eta | \mathbf{q}^2) \vartheta_4(v - \eta | \mathbf{q}^2) \vartheta_4(v + \eta | \mathbf{q}^2), \\ d &= \rho \vartheta_1(2\eta | \mathbf{q}^2) \vartheta_1(v - \eta | \mathbf{q}^2) \vartheta_1(v + \eta | \mathbf{q}^2), \end{aligned} \quad (\text{A.13})$$

where we use the standard theta-functions [6]

$$\vartheta_i(v | \mathbf{q}), \quad i = 1, \dots, 4, \quad \mathbf{q} = e^{i\pi\tau}, \quad \text{Im } \tau > 0, \quad (\text{A.14})$$

with the periods π and $\pi\tau$. Apart from the simple change in the notation for the theta-functions the above parametrization is the same as that given by Eq.(8) of [1]. The theta-functions $\mathbf{H}(z)$ and $\Theta(z)$ of the nome q_B therein are given by

$$q_B = \mathbf{q}^2, \quad \mathbf{H}(z) = \vartheta_1\left(\frac{\pi z}{2\mathbf{K}_B} | \mathbf{q}^2\right), \quad \Theta(z) = \vartheta_4\left(\frac{\pi z}{2\mathbf{K}_B} | \mathbf{q}^2\right), \quad (\text{A.15})$$

where $\mathbf{K}_B$ is the complete elliptic integral of the first kind of the modulus

$$k = \frac{\vartheta_2(0 | \mathbf{q}_B)}{\vartheta_3(0 | \mathbf{q}_B)}. \quad (\text{A.16})$$

The variables v, η and ρ used in [1] (hereafter denoted as v_B, η_B and ρ_B) are related to those in (A.13) by

$$v = \frac{\pi v_B}{2\mathbf{K}_B}, \quad \eta = \frac{\pi \eta_B}{2\mathbf{K}_B}, \quad \rho = \rho_B. \quad (\text{A.17})$$

With these definitions the transfer matrix (A.9) is the same as that given by Eqs.(6)-(8) of ref [1]. We will denote it as $\mathbf{T}^B(v)$, remembering that our variable v is related to Baxter's variable v_B by (A.17).

Throughout this paper we fix the normalization factor ρ in (A.13) as

$$\rho = 2 \vartheta_2(0 | \mathbf{q})^{-1} \vartheta_4(0 | \mathbf{q}^2)^{-1}. \quad (\text{A.18})$$

Note, that the variables (A.5) can be then written as

$$w_i = \frac{1}{2} \vartheta_1(2\eta | \mathbf{q}) \frac{\vartheta_{5-i}(u | \mathbf{q})}{\vartheta_{5-i}(\eta | \mathbf{q})}, \quad i = 1, \dots, 4. \quad (\text{A.19})$$

We shall also make use of two invariants

$$\zeta = \frac{cd}{ab} = \left[\frac{\vartheta_1(2\eta | \mathbf{q}^2)}{\vartheta_4(2\eta | \mathbf{q}^2)} \right]^2, \quad \gamma = \frac{w_3^2 - w_2^2}{w_4^2 - w_1^2} = - \left[\frac{\theta_1(\eta | \mathbf{q}) \theta_4(\eta | \mathbf{q})}{\theta_2(\eta | \mathbf{q}) \theta_3(\eta | \mathbf{q})} \right]^2. \quad (\text{A.20})$$

The transfer matrix $\mathbf{T}^B(v)$ commutes [81, 82] with the Hamiltonian of the XYZ -model (remind that we are assuming the periodic boundary condition),

$$\mathbf{H}_{XYZ} = -\frac{1}{2} \sum_{j=1}^N (J_x \sigma_1^{(j)} \sigma_1^{(j+1)} + J_y \sigma_2^{(j)} \sigma_2^{(j+1)} + J_z \sigma_3^{(j)} \sigma_3^{(j+1)}) \quad (\text{A.21})$$

provided

$$J_x : J_y : J_z = \frac{\vartheta_4(2\eta | \mathbf{q})}{\vartheta_4(0 | \mathbf{q})} : \frac{\vartheta_3(2\eta | \mathbf{q})}{\vartheta_3(0 | \mathbf{q})} : \frac{\vartheta_2(2\eta | \mathbf{q})}{\vartheta_2(0 | \mathbf{q})} . \quad (\text{A.22})$$

Following [82] let us show that this Hamiltonian is simply related to the logarithmic derivative of the transfer matrix at $v = \eta$. It follows from (A.3) and (A.19) that

$$\vartheta_1(2\eta | \mathbf{q}) \left[\frac{d}{dv} \log \mathbf{T}^B(v) \right]_{v=\eta} = N p'_4 \mathbf{I} + \sum_{j=1}^N (p'_1 \sigma_1^{(j)} \sigma_1^{(j+1)} + p'_2 \sigma_2^{(j)} \sigma_2^{(j+1)} + p'_3 \sigma_3^{(j)} \sigma_3^{(j+1)}) , \quad (\text{A.23})$$

where $\mathbf{I}$ denotes the unit operator. Here we used new variables

$$p_1 = \frac{1}{2}(w_1 - w_2 - w_3 + w_4), \quad p_2 = \frac{1}{2}(-w_1 + w_2 - w_3 + w_4) , \quad (\text{A.24})$$

$$p_3 = \frac{1}{2}(-w_1 - w_2 + w_3 + w_4), \quad p_4 = \frac{1}{2}(w_1 + w_2 + w_3 + w_4) . \quad (\text{A.25})$$

Their v -derivatives evaluated at $v = \eta$ denoted as

$$p'_i = \left. \frac{dp_i}{dv} \right|_{v=\eta} . \quad (\text{A.26})$$

Note that

$$p_1 + p_2 + p_3 + p_4 = 2w_4 . \quad (\text{A.27})$$

Using (A.19) one can readily show that

$$2p'_i = \vartheta'_1(0 | \mathbf{q}) \frac{\vartheta_{5-i}(2\eta | \mathbf{q})}{\vartheta_{5-i}(0 | \mathbf{q})}, \quad i = 1, 2, 3 . \quad (\text{A.28})$$

Combining the last relation with (A.23) one can express the Hamiltonian (A.21) as

$$\mathbf{H}_{XYZ} = -\frac{J_x \vartheta_4(0 | \mathbf{q})}{\vartheta_4(2\eta | \mathbf{q}) \vartheta'_1(0 | \mathbf{q})} \left\{ \vartheta_1(2\eta | \mathbf{q}) \frac{d}{dv} \log \mathbf{T}^B(v) - N p'_4 \mathbf{I} \right\} \Big|_{v=\eta} \quad (\text{A.29})$$

In the main text of the paper we mostly use the spectral parameter u which is related to variable v in (A.13) as

$$u = v - \pi\tau/2 . \quad (\text{A.30})$$

In (2.64) we also define the renormalized transfer-matrix $\mathbf{T}(u)$ differing from $\mathbf{T}^B(u + \pi\tau/2)$ by a simple u -dependent factor. Of course, this factor could have been included in the normalization of the Boltzmann weights (A.13), but we prefer not to do so and keep a clear distinction between two alternative normalizations and, in fact, to use both of them. In particular, the Baxter's normalization and the variable v are more suitable in writing explicit expressions for the eigenvalues of the transfer matrix presented in Sect. 4.

There are two related, but different, constructions of the $\mathbf{Q}$ -matrix for the 8V-model, given in Baxter's 1972 and 1973 papers [1] and [2]. The results of [1] apply for rational η 's and an arbitrary number of sites N , while the those of [2] apply to arbitrary η 's and even N . We will denote the corresponding $\mathbf{Q}$ -matrices as $\mathbf{Q}^{(72)}(v)$ and $\mathbf{Q}^{(73)}(v)$. They obey the same TQ-equation (Eq.(4.2) of [1] and Eq.(87) of [2]),

$$\mathbf{T}^B(v) \mathbf{Q}^B(v) = \phi(v - \eta) \mathbf{Q}^B(v + 2\eta) + \phi(v + \eta) \mathbf{Q}^B(v - 2\eta), \quad (\text{A.31})$$

where

$$\phi^B(v) = (\vartheta_1(v|q))^N. \quad (\text{A.32})$$

and possess the same periodicity properties

$$\mathbf{Q}^B(v + \pi) = \mathcal{S} \mathbf{Q}^B(v), \quad \mathbf{Q}^B(v + 2\pi\tau) = \mathbf{q}^{-2N} e^{-2ivN} \mathbf{Q}^B(v). \quad (\text{A.33})$$

where $\mathcal{S}$ is defined in (A.12) and $\mathbf{Q}^B(v)$ stands for either of $\mathbf{Q}^{(72)}(v)$ or $\mathbf{Q}^{(73)}(v)$. Nevertheless, as noticed in [14], these two $\mathbf{Q}$ -matrices do not coincide. Of course, they can only be compared for rational η and even N when both of them can be constructed. As shown in [14] the matrices $\mathbf{Q}^{(72)}(v)$ and $\mathbf{Q}^{(73)}(v)$ have different eigenvalues for degenerate eigenstates of the transfer matrix. This phenomenon is obviously related with the non-uniqueness of the eigenvectors for the degenerate eigenstates [13] and demonstrates the impossibility of an universal definition of the $\mathbf{Q}$ -matrix in the zero-field 8V-model at rational η (cf. [54]).

The difference between $\mathbf{Q}^{(72)}(v)$ and $\mathbf{Q}^{(73)}(v)$ can be traced back [14] to their commutativity properties with the operator $\mathcal{R}$. The matrix $\mathbf{Q}^{(73)}(v)$ (defined for even N only) always commutes with $\mathcal{R}$. The matrix $\mathbf{Q}^{(72)}(v)$ (defined for all values of N) does not commute with $\mathcal{R}$, irrespectively to whether N is odd or even. On the other hand both these matrices commute with $\mathcal{S}$, so that the first equation immediately translates into the (real period) periodicity for the eigenvalues (the first relation in (2.66)). The second equation in (A.33) implies only the $2\pi\tau$ -periodicity relation (the second relation in (2.66)).

To complete our discussion of various constructions for the $\mathbf{Q}$ -matrix, note that in [5] Baxter considered a modified version of his $\mathbf{Q}^{(73)}$ matrix. There he used yet another set of parameters, namely the variables v and λ , which we denote here as $v^{(82)}$ and $\lambda^{(82)}$. They are related to v and η in (A.13) as

$$v^{(82)} = \frac{2iK_B}{\pi}(2v - \pi), \quad \lambda^{(82)} = \frac{2iK_B}{\pi}(\pi - 2\eta) \quad (\text{A.34})$$

where K_B is the same as in (A.17). Writing the $\mathbf{Q}$ -matrix of [5] as $\mathbf{Q}^{(82)}(v^{(82)}, \lambda^{(82)})$ and that of [2] as $\mathbf{Q}^{(73)}(v, \eta)$, and using the explanations in Sect.7 of [13] it is not difficult to verify that

$$\mathbf{Q}^{(73)}(v, \eta) = \mathcal{R} \mathcal{D} \mathbf{Q}^{(82)}(v^{(82)}, \lambda^{(82)}) \quad (\text{A.35})$$

where the operator

$$\mathcal{D} = e^{+\frac{i\pi}{4}\sigma_3} \otimes e^{-\frac{i\pi}{4}\sigma_3} \otimes e^{+\frac{i\pi}{4}\sigma_3} \otimes \dots \otimes e^{-\frac{i\pi}{4}\sigma_3} \quad (\text{A.36})$$

is defined similarly to those in (A.12). Using the relations

$$\mathcal{R} \mathcal{D} = (-1)^{N/2} \mathcal{D} \mathcal{R} \mathcal{S}, \quad \mathbf{Q}^{(73)}(v + \pi\tau) = \mathbf{q}^{-N/2} e^{-ivN} \mathcal{R} \mathbf{Q}^{(73)}(v), \quad (\text{A.37})$$

one can show that

$$\mathbf{Q}^{(82)}(v^{(82)} + 2K'_B) = (q_B)^{-N/4} \exp(\pi v^{(82)} N / 4K_B) \mathcal{R} \mathcal{S} \mathbf{Q}^{(82)}(v^{(82)}) \quad (\text{A.38})$$

which is exactly Eq.(10.5.43a) of [5].

The partition function (A.2) can be regarded as a function of four parameters w_1, w_2, w_3, w_4 , defined in (A.5). So one can write it as $Z(w_1, w_2, w_3, w_4)$. When both M and N are even, it possesses the symmetry

$$Z(w_1, w_2, w_3, w_4) = Z(\pm w_i, \pm w_j, \pm w_k, \pm w_l), \quad (\text{A.39})$$

where $\{i, j, k, l\}$ is an arbitrary permutation of $\{1, 2, 3, 4\}$ and all signs can be chosen independently. Using this symmetry one can always rearrange w_i so that

$$w_1 > w_2 > w_3 > |w_4|. \quad (\text{A.40})$$

The partition function-per-site, $\kappa(v)$, is defined as

$$\log \kappa(v) = \lim_{M, N \rightarrow \infty} \frac{1}{MN} \log Z = \lim_{N \rightarrow \infty} \frac{1}{N} \log \Lambda_0^B(v) \quad (\text{A.41})$$

where $\Lambda_0^B(v)$ is the maximum eigenvalue of the transfer-matrix $\mathbf{T}^B(v)$. In the (unphysical) regime (A.40) it reads [1]

$$\log \kappa(v) = \log c + 2 \sum_{n=1}^{\infty} \frac{\sinh^2((\tilde{\tau}_B - l_B)n) (\cosh(nl_B) - \cosh(n\alpha_B))}{n \sinh(2n\tilde{\tau}_B) \cosh(nl_B)}, \quad (\text{A.42})$$

where

$$\alpha_B = -2iv, \quad l_B = -2i\eta, \quad \tilde{\tau}_B = -\pi i \tau. \quad (\text{A.43})$$

In this paper we consider the disordered regime¹⁹

$$0 < \eta < \pi/2, \quad \eta < v < \pi - \eta, \quad (\text{A.44})$$

which corresponds to a different ordering of the variables w_i , namely,

$$w_4 > w_1 > w_2 > w_3 > 0. \quad (\text{A.45})$$

This regime can be mapped into (A.40) by the following transformation of the weights

$$a' = \frac{a - b + c - d}{2}, \quad b' = \frac{a - b - c + d}{2}, \quad c' = \frac{a + b + c + d}{2}, \quad d' = \frac{a + b - c - d}{2}, \quad (\text{A.46})$$

which is equivalent to

$$w'_1 = w_4, \quad w'_2 = w_1, \quad w'_3 = w_2, \quad w'_4 = w_3, \quad (\text{A.47})$$

where w'_1, w'_2, w'_3, w'_4 are defined by (A.5) with a, b, c, d replaced by a', b', c', d' . It is easy to check that if w_1, w_2, w_3, w_4 belong to the regime (A.45) then w'_1, w'_2, w'_3, w'_4 belong to (A.40). Putting (A.13) into (A.46) it is not very difficult to show that the “transformed” weights a', b', c', d' can be parameterized by same formulae (A.13) provided one makes the following substitution

$$v \rightarrow \frac{v - \pi/2}{\tau}, \quad \eta \rightarrow \frac{\eta - \pi/2}{\tau}, \quad \tau \rightarrow -\frac{1}{\tau}, \quad \rho \rightarrow \rho_1 \rho, \quad (\text{A.48})$$

where

$$\rho_1 = -i(-i\tau)^{-1/2} \exp\left\{-\frac{i}{\pi\tau}[v(v - \pi) + 3\eta(\eta - \pi) + \pi^2]\right\}. \quad (\text{A.49})$$

¹⁹See [5] for the classification of the regimes of the 8V-model. Beware that the variables w_i therein are numbered differently as compared to this work. Here we follow the original papers [1–4].

Thus, to find the partition function-per-site in the regime (A.45) one needs to substitute parameters α_B, l_B and τ_B in (A.42) by

$$\alpha_B = \frac{i(\pi - 2v)}{\tau}, \quad l_B = \frac{i(\pi - 2\eta)}{\tau}, \quad \tilde{\tau}_B = \frac{i\pi}{\tau}, \quad (\text{A.50})$$

and replace c by c' . After straightforward calculations one obtains

$$\log \kappa(v) = \log \vartheta_1(v + \eta | \mathbf{q}) + \frac{2i\eta(v - \eta)}{\pi\tau} + \sum_{n=1}^{\infty} \frac{\sinh(2\eta x_n) \sinh(2(v - \eta) x_n)}{n \sinh(\pi x_n) \cosh((\pi - 2\eta) x_n)}, \quad (\text{A.51})$$

where $x_n = in/\tau$. The above formula can be equivalently rewritten as

$$\log \kappa(v) = \log \vartheta_1(v | \mathbf{q}) + \frac{i\eta(\pi - 3\eta)}{\pi\tau} + 2 \sum_{n=1}^{\infty} \frac{\sinh(\eta x_n) \sinh((\pi - 3\eta) x_n) \cosh((\pi - 2v) x_n)}{n \sinh(\pi x_n) \cosh((\pi - 2\eta) x_n)}. \quad (\text{A.52})$$

Note that for $\eta = \pi/3$ the second and third terms in the last expression vanish, so it simplifies to

$$\log \kappa(v) = \log \vartheta_1(v | \mathbf{q}), \quad \eta = \pi/3. \quad (\text{A.53})$$

Recall that the above derivation was based on the symmetry (A.39) valid for even values of N only. However, since in the disordered regime (A.45) the Boltzmann weights (A.13) are strictly positive, the result (A.53), obviously, does not depend on the way the limit $N \rightarrow \infty$ is taken. Eqs. (A.41) and (A.53) imply that the asymptotics on the largest eigenvalue of $\mathbf{T}^B(v)$ in this special case has an extremely simple form $\Lambda_0^B(v) \simeq (\vartheta_1(v | \mathbf{q}))^N$. Remarkably, as it was argued in [16, 42, 86], for odd values of N this asymptotics is *exact* even for a finite lattice,

$$\Lambda_0^B(v) = (a + b)^N = \vartheta_1^N(v | q), \quad N = 2n + 1, \quad \eta = \pi/3. \quad (\text{A.54})$$

We have also checked this fact analytically up to $N = 15$.

References

- [1] Baxter, R. J. Partition function of the eight-vertex lattice model. *Ann. Physics* **70** (1972) 193–228.
- [2] Baxter, R. J. Eight-vertex model in lattice statistics and one-dimensional anisotropic Heisenberg chain. I. Some fundamental eigenvectors. *Ann. Phys.* **76** (1973) 1–24.
- [3] Baxter, R. J. Eight-vertex model in lattice statistics and one-dimensional anisotropic Heisenberg chain. II. Equivalence to a generalized Ice-type lattice model. *Ann. Phys.* **76** (1973) 25–47.
- [4] Baxter, R. J. Eight-vertex model in lattice statistics and one-dimensional anisotropic Heisenberg chain. III. Eigenvectors of the transfer matrix and Hamiltonian. *Ann. Phys.* **76** (1973) 48–71.
- [5] Baxter, R. J. *Exactly Solved Models in Statistical Mechanics*. Academic Press Inc., London, 1982.
- [6] Whittaker, E. and Watson, G. *A course of modern analysis*. “Cambridge University Press”, Cambridge, 1996.

- [7] Faddeev, L. D., Sklyanin, E. K., and Takhtajan, L. A. The Quantum Inverse Problem Method. 1. Theor. Math. Phys. **40** (1980) 688.
- [8] Bazhanov, V. V., Lukyanov, S. L., and Zamolodchikov, A. B. Integrable structure of conformal field theory. II. Q -operator and DDV equation. Comm. Math. Phys. **190** (1997) 247–278. [[hep-th/9604044](#)].
- [9] Bazhanov, V. V., Lukyanov, S. L., and Zamolodchikov, A. B. Integrable structure of conformal field theory. III. The Yang-Baxter Relation. Comm. Math. Phys. **200** (1999) 297–324. [arXiv:hep-th/9805008](#).
- [10] Krichever, I., Lipan, O., Wiegmann, P., and Zabrodin, A. Quantum integrable models and discrete classical Hirota equations. Comm. Math. Phys. **188** (1997) 267–304. [[hep-th/9604080](#)].
- [11] Takhtajan, L. A. and Faddeev, L. D. The quantum method for the inverse problem and the XYZ Heisenberg model. Uspekhi Mat. Nauk **34(5)** (1979) 13–63. (English translation: Russian Math. Surveys **34(5)** (1979) 11–68).
- [12] Felder, G. and Varchenko, A. Algebraic Bethe ansatz for the elliptic quantum group $E_{\tau,\eta}(\mathfrak{sl}_2)$. Nuclear Phys. B **480** (1996) 485–503.
- [13] Baxter, R. J. Completeness of the Bethe ansatz for the six- and eight-vertex models. J. Statist. Phys. **108** (2002) 1–48.
- [14] Fabricius, K. and McCoy, B. M. New developments in the eight vertex model. J. Statist. Phys. **111** (2003) 323–337.
- [15] Fabricius, K. and McCoy, B. M. Bethe’s equation is incomplete for the XXZ model at roots of unity. J. Statist. Phys. **103** (2001) 647–678.
- [16] Bazhanov, V. V. and Mangazeev, V. V. Eight-vertex model and non-stationary Lamé equation. J. Phys. A **38** (2005) L145–153. [arXiv/hep-th/0411094](#).
- [17] Bazhanov, V. V. and Mangazeev, V. V. Eight vertex model and Painlevé VI. J. Phys. A **39** (2006) 12235–12243. [arXiv:hep-th/0602122](#).
- [18] Bazhanov, V. V. and Mangazeev, V. V. Analytic theory of the eight vertex model II. Partial differential equations and Painlevé transcendents. In preparation.
- [19] Baxter, R. J. Generalized ferroelectric model on a square lattice. Stud. Appl. Math. **1** (1971) 51–69.
- [20] Andrews, G. E., Baxter, R. J., and Forrester, P. J. 8-vertex SOS model and generalized Rogers-Ramanujan-type identities. J. Stat. Phys. **35** (1984) 193–266.
- [21] Bazhanov, V. V., Lukyanov, S. L., and Zamolodchikov, A. B. Quantum field theories in finite volume: excited state energies. Nuclear Phys. B **489** (1997) 487–531. [[hep-th/9607099](#)].
- [22] Kulish, P. P., Reshetikhin, N. Y., and Sklyanin, E. K. Yang-Baxter equations and representation theory. I. Lett. Math. Phys. **5** (1981) 393–403.
- [23] Bazhanov, V. V., Lukyanov, S. L., and Zamolodchikov, A. B. 1997, unpublished.

- [24] Fendley, P., Ludwig, A. W. W., and Saleur, H. Exact non-equilibrium transport through point contacts in quantum wires and fractional quantum Hall devices. *Phys. Rev. B* **52** (1995) 8934–8950. [[cond-mat/9503172](#)].
- [25] Baxter, R. J. and Pearce, P. A. Hard hexagons: interfacial tension and correlation length. *J. Phys. A* **15** (1982) 897–910.
- [26] Bazhanov, V. V. and Reshetikhin, N. Y. Critical RSOS models and conformal field theory. *Internat. J. Modern Phys. A* **4** (1989) 115–142.
- [27] Date, E., Jimbo, M., Miwa, T., and Okado, M. Fusion of the eight vertex SOS model. *Lett. Math. Phys.* **12** (1986) 209–215. Errata: *ibid* **14** (1987) 97.
- [28] Fabricius, K. and McCoy, B. M. New developments in the eight vertex model. II. Chains of odd length. *J. Stat. Phys.* **120** (2005) 37–70.
- [29] Bazhanov, V. V., Lukyanov, S. L., and Zamolodchikov, A. B. Integrable structure of conformal field theory, quantum KdV theory and thermodynamic Bethe Ansatz. *Comm. Math. Phys.* **177** (1996) 381–398. [[hep-th/9412229](#)].
- [30] Stroganov, Y. G. A new calculation method for partition functions in some lattice models. *Phys. Lett. A* **74** (1979) 116–118.
- [31] Zamolodchikov, A. B. Z_4 -symmetric factorized S -matrix in two space-time dimensions. *Comm. Math. Phys.* **69** (1979) 165–178.
- [32] Baxter, R. J. The inversion relation method for some two-dimensional exactly solved models in lattice statistics. *J. Statist. Phys.* **28** (1982) 1–41.
- [33] Takahashi, M. Simplification of thermodynamic Bethe-ansatz equations. [arXiv.org:cond-mat/0010486](#), 2000.
- [34] Stroganov, Y. G. 1979, unpublished.
- [35] Lieb, E. H. Residual Entropy of Square Ice. *Phys. Rev.* **162** (1967) 162–172.
- [36] Reshetikhin, N. Y. The functional equation method in the theory of exactly soluble quantum systems. *Zh. Èksper. Teoret. Fiz.* **84** (1983) 1190–1201.
- [37] Kirillov, A. N. and Reshetikhin, N. Y. Exact solution of the integrable XXZ Heisenberg model with arbitrary spin. I. The ground state and the excitation spectrum. *J. Phys. A* **20** (1987) 1565–1585.
- [38] Sklyanin, E. K. Some algebraic structures connected with the Yang-Baxter equation. *Func. Anal. Appl.* **16** (1982) 263–270.
- [39] Sklyanin, E. K. Some algebraic structures connected with the Yang-Baxter equation. Representations of a quantum algebra. *Func. Anal. Appl.* **17** (1983) 273–284.
- [40] Krichever, I. and Zabrodin, A. Vacuum curves of elliptic L -operators and representations of the Sklyanin algebra. In *Moscow Seminar in Mathematical Physics*, volume 191 of *Amer. Math. Soc. Transl. Ser. 2*, pages 199–221. Amer. Math. Soc., Providence, RI, 1999.
- [41] Fabricius, K. and McCoy, B. M. Functional equations and fusion matrices for the eight vertex model. *Publ. Res. Inst. Math. Sci.* **40** (2004) 905–932.

- [42] Baxter, R. J. Solving models in statistical mechanics. *Adv. Stud. Pure Math.* **19** (1989) 95–116.
- [43] Baxter, R. J. Hard hexagons: exact solution. *J. Phys. A* **13** (1980) L61–L70.
- [44] Nepomechie, R. I. Functional relations and Bethe Ansatz for the XXZ chain. *Journal of Statistical Physics* **111** (2003) 1363.
- [45] Fendley, P. Airy functions in the thermodynamic Bethe ansatz. *Lett. Math. Phys.* **49** (1999) 229–233.
- [46] Stroganov, Y. G. The XXZ-spin chain with the asymmetry parameter $\Delta = -1/2$. Computation of the simplest correlators. *Teoret. Mat. Fiz.* **129** (2001) 345–359.
- [47] Yang, C. N. and Yang, C. P. Thermodynamics of a one-dimensional system of bosons with repulsive delta-function interaction. *J. Mathematical Phys.* **10** (1969) 1115–1122.
- [48] Takahashi, M. and Suzuki, M. One-Dimensional Anisotropic Heisenberg Model at Finite Temperatures. *Progr. Theor. Phys.* **48** (1972) 2187–2209.
- [49] Zamolodchikov, A. B. On the thermodynamic Bethe Ansatz equations for reflectionless *ADE* scattering theories. *Phys. Lett. B* **253** (1991) 391–394.
- [50] Klümper, A. and Pearce, P. A. Conformal Weights of RSOS Lattice Models and their Fusion Hierarchies. *Phys. A* **183** (1992) 304–350.
- [51] Kuniba, A., Nakanishi, T., and Suzuki, J. Functional relations in solvable lattice models. I. Functional relations and representation theory. *Internat. J. Modern Phys. A* **9** (1994) 5215–5266. [[hep-th/9309137](#)].
- [52] Fendley, P., Lesage, F., and Saleur, H. Solving 1D plasmas and 2D boundary problems using Jack polynomials and functional relations. *J. Statist. Phys.* **79** (1995) 799–819.
- [53] Dorey, P. and Tateo, R. Excited states by analytic continuation of TBA equations. *Nuclear Phys. B* **482** (1996) 639–659.
- [54] Bazhanov, V. V. and Stroganov, Y. G. Chiral Potts model as a descendant of the six-vertex model. *J. Statist. Phys.* **59** (1990) 799–817.
- [55] von Gehlen, G. and Rittenberg, V. $Z(n)$ -symmetric quantum chains with an infinite set of conserved charges and $Z(n)$ zero modes. *Nucl. Phys.* **B257** (1985) 351.
- [56] Au-Yang, H., McCoy, B. M., Perk, J. H. H., Tang, S., and Yan, M.-L. Commuting transfer matrices in the chiral Potts models: Solutions of Star triangle equations with genus > 1 . *Phys. Lett.* **A123** (1987) 219–223.
- [57] Baxter, R. J., Perk, J. H. H., and Au-Yang, H. New solutions of the star triangle relations for the chiral Potts model. *Phys. Lett.* **A128** (1988) 138–142.
- [58] Kashiwara, M. and Miwa, T. A class of elliptic solutions to the star-triangle relation. *Nuclear Physics B* **275** (1986) 121–134.
- [59] Hasegawa, K. and Yamada, Y. Algebraic derivation of the broken Z_N -symmetric model. *Phys. Lett. A* **146** (1990) 387–396.

- [60] Gaudin, M. La relation étoile-triangle d'un modèle elliptique $\mathbf{Z}_N$. J. Physique I **1** (1991) 351–361.
- [61] Pasquier, V. and Gaudin, M. The periodic Toda chain and a matrix generalization of the Bessel function recursion relations. J. Phys. A **25** (1992) 5243–5252.
- [62] Derkachov, S. É. Baxter's Q -operator for the homogeneous XXX spin chain. J. Phys. A **32** (1999) 5299–5316.
- [63] Sklyanin, E. K. Bäcklund transformations and Baxter's Q -operator. In *Integrable systems: from classical to quantum (Montréal, QC, 1999)*, volume 26 of *CRM Proc. Lecture Notes*, pages 227–250. Amer. Math. Soc., Providence, RI, 2000.
- [64] Deguchi, T., Fabricius, K., and McCoy, B. M. The sl_2 loop algebra symmetry of the six-vertex model at roots of unity. J. Stat. Phys. **102** (2001) 701–736.
- [65] Kashaev, R. M. The non-compact quantum dilogarithm and the Baxter equations. J. Statist. Phys. **102** (2001) 923–936.
- [66] Fabricius, K. and McCoy, B. M. Evaluation Parameters and Bethe Roots for the Six-Vertex Model at Roots of Unity. In *MathPhys Odyssey 2001 (Hardcover)*, ed. M. Kashiwara and T. Miwa, volume 23 of *Progress in Mathematical Physics*, pages 119–144, Boston, 2002. Birkhäuser.
- [67] Smirnov, F. A. Dual Baxter equations and quantization of affine Jacobian. [[math-ph/0001032](#)].
- [68] Zabrodin, A. Commuting difference operators with elliptic coefficients from Baxter's vacuum vectors. J. Phys. A **33** (2000) 3825–3850.
- [69] Fabricius, K. and McCoy, B. M. An elliptic current operator for the eight-vertex model. J. Phys. A: Math. Gen. **39** (2006) 14869–14886.
- [70] Khoroshkin, S. M., Stolin, A. A., and Tolstoy, V. N. Generalized Gauss decomposition of Trigonometric R -matrices. Modern Phys. Lett. A **10** (1995) 1375–1392.
- [71] A. A. Belavin, A. V. Odesskii, R. A. U. New relations in the algebra of the Baxter Q -operators. Theor. Math. Phys. **130** (2002) 323–350.
- [72] Korff, C. Auxiliary matrices on both sides of the equator. J. Phys. A: Math. Gen. **38** (2005) 47–67.
- [73] Korff, C. A Q -operator identity for the correlation functions of the infinite XXZ spin-chain. J. Phys. A: Math. Gen. **38** (2005) 6641–6658.
- [74] Boos, H., Jimbo, M., Miwa, T., Smirnov, F., and Takeyama, Y. Hidden Grassmann structure in the XXZ model. [hep-th/0606280](#).
- [75] Albertini, G., McCoy, B. M., and Perk, J. H. H. Eigenvalue spectrum of the superintegrable chiral Potts model. In *Integrable systems in quantum field theory and statistical mechanics*, volume 19 of *Adv. Stud. Pure Math.*, pages 1–55. Academic Press, Boston, MA, 1989.
- [76] Baxter, R. J., Bazhanov, V. V., and Perk, J. H. H. Functional relations for transfer matrices of the chiral Potts model. Internat. J. Modern Phys. B **4** (1990) 803–870.

- [77] Pronko, G. P. and Stroganov, Y. G. Bethe equations ‘on the wrong side of the equator’. J. Phys. A **32** (1999) 2333–2340.
- [78] Deguchi, T. Construction of some missing eigenvectors of the XYZ spin chain at the discrete coupling constants and the exponentially large spectral degeneracy of the transfer matrix. J. Phys. A **35** (2002) 879–895.
- [79] Johnson, J. D., Krinsky, S., and McCoy, B. M. Vertical-Arrow Correlation Length in the Eight-Vertex Model and the Low-Lying Excitations of the X-Y-Z Hamiltonian. Phys. Rev. A **8** (1973) 2526–2547.
- [80] Bazhanov, V. V., Lukyanov, S. L., and Zamolodchikov, A. B. Higher-level eigenvalues of Q-operators and Schroedinger equation. Adv. Theor. Math. Phys. **7** (2003) 711–725.
- [81] Sutherland, B. Two-Dimensional Hydrogen Bonded Crystals without the Ice Rule. Journal of Mathematical Physics **11** (1970) 3183–3186.
- [82] Baxter, R. J. One-dimensional anisotropic Heisenberg chain. Ann. Phys. **70** (1972) 323–337.
- [83] Batchelor, M. T., Barber, M. N., and Pearce, P. A. Bethe ansatz calculations for the eight-vertex model on a finite strip. J. Statist. Phys. **49** (1987) 1117–1163.
- [84] Sutherland, B. Low-Lying Eigenstates of the One-Dimensional Heisenberg Ferromagnet for any Magnetization and Momentum. Phys. Rev. Lett. **74** (1995) 816.
- [85] Beisert, N., Staudacher, M., Minahan, J. A., and Zarembo, K. Stringing spins and spinning strings. J. High Energy Phys. (2003) 010, 28 pp. (electronic).
- [86] Stroganov, Y. The 8-vertex model with a special value of the crossing parameter and the related XYZ spin chain. In *Integrable structures of exactly solvable two-dimensional models of quantum field theory (Kiev, 2000)*, volume 35 of *NATO Sci. Ser. II Math. Phys. Chem.*, pages 315–319. Kluwer Acad. Publ., Dordrecht, 2001.